



Consent for COVID-19 Immunization

For use at Alberta Health Services (AHS) immunization programs. Use this form when a parent or alternate decision-maker is **not** able to be with the person being immunized at an AHS immunization service. An alternate decision-maker could be an agent, guardian, specific decision-maker or co-decision-maker. The parent or alternate decision-maker should **complete this form** and **send it** with the person being immunized, to the AHS immunization service. For alternate decision-maker – please also send a copy of documents to show that you are authorized to be the alternate decision-maker.

Personal information for the person being immunized					
Name (Last, First, Middle)	Date of Birth (dd-Mon-yyyy)				
Personal Health Number (PHN)	Gender				
Health information for the person being immunized (If you need more space, use the other side of this form.)					
Does this person have any allergies, including allergies to any vaccine, medicine, or for If yes, describe	ood? □ No □ Yes				
Does this person have any chronic illness? (<i>List all if more than one</i>) If yes, describe	□ No □ Yes				
Has this person ever had Capillary Leak Syndrome?	🗆 No 🛛 Yes				
Is this person taking any medicine?					
Is this person pregnant?	ing? 🗆 No 🗆 Yes				
Has this person had COVID-19 vaccine before?					
Has this person ever had a side effect from COVID-19 immunization? If yes, describe	□ No □ Yes				
Will this person get another vaccine in the 14 days before they get the COVID-19 vaccine?					
Consent					
 I confirm that I have read the attached COVID-19 vaccine information. I know about and understand the risks, benefits, and common side effects of this vaccine. Any questions I may have had about this person getting this vaccine have been answered by calling the local public health office or Health Link at 811. I understand the information I have been given. I understand this consent is for all doses of the vaccine. I will contact the local public health office or the healthcare provider giving the COVID-19 vaccine if this person: has any changes to their health before getting any dose of the COVID-19 vaccine gets another vaccine in the 14 days before they get any dose of the COVID-19 vaccine has a severe or unusual side effect after any dose of the COVID-19 vaccine (other than the expected side effects listed on the COVID-19 vaccine (for ages 12 years and older) I consent to this person getting the (only select one): Pfizer-BioNTech mRNA COVID-19 vaccine (for ages 12 years and older) Moderna mRNA COVID-19 vaccine (for ages 12 years and older) Either the Pfizer-BioNTech or Moderna mRNA COVID-19 vaccine (for ages 18 years and older) AstraZeneca/COVISHIELD viral vector-based COVID-19 vaccine (for ages 18 years and older) I understand that I may withdraw this consent at any time by calling the local public health office or healthcare provider giving the COVID-19 vaccine. 					
Printed name of person giving consent Daytime phone	Other phone				
Relationship to person Parent (with legal authority to consent) Co-decision-maker Signature of person giving consent	□ Agent Date <i>(dd-Mon-yyyy)</i>				

Alberta Health Services collects health information according to Section 20 of the Health Information Act (HIA). This information is used to provide health services, determine eligibility for health services, or to carry out any other purpose authorized by the HIA. If you have any questions about this, please ask the healthcare provider giving the immunization or contact your local public health office. If you do not know your local public health office, call Health Link at 811 to get this information.



For Office Use Only

Name (Last, First, Middle)	PHN				
Telephone/Fax Consent					
Mode by which consent was received	□ Fax/Scan	□ Telephone			
Name of healthcare provider obtaining the cons	sent	Date (dd-Mon-yyyy) Time			
Signature of healthcare provider obtaining the consent					
Consent Using an Interpreter (for non-English speaking parent/client)					
Interpreter's name or ID #	Phone Date (dd-		Mon-yyyy)	Time	
Notes (For Office Use Only)				1	



COVID-19 vaccines

Immunization protects you from disease. Get protected, get immunized.

- Vaccines make your immune system stronger. They build antibodies to help prevent diseases.
- Immunization is safe. It is much safer to get immunized than to get this disease.

What are COVID-19 vaccines?

COVID-19 vaccines protect against the SARS-CoV-2 virus (also known as COVID-19). The virus causes an infection in the lungs and airways and is a type of respiratory illness. Go to <u>ahs.ca/covid</u> to learn more about COVID-19. There are 2 types of COVID-19 vaccines approved for use in Canada:

- **mRNA vaccines:** The Pfizer-BioNTech and Moderna vaccines are mRNA vaccines.
- Viral vector-based vaccines: The AstraZeneca/COVISHIELD vaccine is a viral vector-based vaccine.

Learn more about COVID-19 vaccines at <u>canada.ca/en/health-canada/services/drugs-health-products/covid19-industry/drugs-vaccines-treatments/vaccines.html</u>.

Who should get a COVID-19 vaccine?

You should get a COVID-19 vaccine if you are age 12 years or older. Everyone is at risk of COVID-19. COVID-19 vaccines are free.

If your child is getting a COVID-19 vaccine

Children under age 18 years need a parent or guardian to give consent for their immunization. If a parent or guardian cannot attend the appointment, they can give their written consent. Please complete the consent form found on <u>ahs.ca/VaccineUnder18</u>. In some cases, children under age 18 years may be able to give their own consent.

Can I get the COVID-19 vaccine if I am pregnant or breastfeeding?

Yes, you can get the COVID-19 vaccine if you are pregnant or breastfeeding. When you are pregnant, you have a higher risk of getting very sick from COVID-19. Getting a COVID-19 vaccine during pregnancy lowers your risk of getting seriously ill from COVID-19.

It is best to get an mRNA COVID-19 vaccine if you are pregnant. Research to date shows that it is the safest type of COVID-19 vaccine to get during pregnancy.

If you are pregnant or breastfeeding and have questions about getting the COVID-19 vaccine, talk to your healthcare provider.

How many doses do I need?

Moderna, Pfizer-BioNTech, and AstraZeneca/COVISHIELD vaccines require at least 2 doses.

You may need additional doses of the COVID-19 vaccine to give you more protection if you have a high risk of severe disease, such as you:

- have a health problem that weakens your immune system, such as you have had or will have an organ or stem cell transplant or you have kidney disease and need dialysis
- take certain medicines that weaken your immune system
- live in a specific type of seniors congregate living facility, such as long term care

If you are travelling outside of Canada, you may need additional doses. This only applies if your destination requires you to have a certain COVID-19 vaccine series.

If you need additional doses, your healthcare provider will let you know when you should have them. Visit <u>alberta.ca/covid19-vaccine.aspx</u> for more information.

How well do the COVID-19 vaccines work?

If you are healthy and get all the doses you need, the vaccines give you very good protection against COVID-19. The mRNA vaccines are somewhat more effective than the viral vector-based vaccines.

You need both doses for the greatest and longest lasting protection. All COVID-19 vaccines work very well to lower your risk of getting seriously ill from COVID-19. They also lower your risk of needing to be in the hospital.

At this time, there is not enough information to know how long protection lasts or if everyone needs additional doses. It is still important to practice the recommended public health measures to prevent the spread of COVID-19, even if you have had the vaccine. Please visit <u>ahs.ca/covid</u> for the most up to date recommendations.

Which COVID-19 vaccine will I get?

You will get an mRNA vaccine for your first dose unless you are unable to receive or do not want an mRNA vaccine.

I got an mRNA vaccine as my first dose. Can I get a different mRNA vaccine for my second dose?

Usually you get the same vaccine for your first and second dose. But you can get a different mRNA vaccine for your second dose if your first-dose vaccine is not available or you prefer a different available mRNA vaccine. Both Pfizer-BioNTech and Moderna COVID-19 vaccines work in the same way.

Whichever vaccine you get for your second dose completes your immunization series and protects you against COVID-19.

Go to <u>ahs.ca/seconddose</u> for more information.

What if I got the AstraZeneca/COVISHIELD vaccine as my first dose?

If you got the AstraZeneca/COVISHIELD vaccine for your first dose, you can choose the AstraZeneca/COVISHIELD vaccine or an mRNA vaccine for your second dose.

The AstraZeneca/COVISHIELD vaccine is a safe vaccine that works well to protect you against COVID-19. But some people may choose an mRNA vaccine to complete their COVID-19 immunization.

No matter which vaccine you choose, the second dose offers protection against COVID-19 and will complete your two-dose series.

Go to <u>ahs.ca/seconddose</u> for more information about getting an mRNA vaccine after an AstraZeneca/COVISHIELD vaccine.

If I need additional doses, what vaccine will I get?

You will get an mRNA vaccine for any additional doses. If you are unable to receive an mRNA vaccine, your healthcare provider will let you know which vaccine you can get.

Where can I get a COVID-19 vaccine?

Go to <u>ahs.ca/covidvaccine</u> to find out where and when you can get the COVID-19 vaccine.

Are there side effects from COVID-19 vaccines?

There can be side effects from COVID-19 vaccines, but they tend to be mild and go away in a few days. Side effects may include:

- redness, warmth, swelling, bruising, itching, or feeling sore where you
 had the needle
- feeling tired or unwell
- a headache
- a fever or chills
- body aches or sore joints
- feeling sick to your stomach (nausea), vomiting (throwing up), or loose stool (diarrhea)
- swollen lymph nodes
- a sore throat, cough, or runny nose
- a reduced sense of touch or a feeling of numbness
- feeling dizzy
- pain in your legs or arms

Current information shows that the risk of side effects after additional doses of COVID-19 vaccine is similar to the risk after the first or second dose. Research continues to find out more about the risk of side effects after additional doses of COVID-19 vaccine.

It is important to stay at the clinic for 15 minutes after your vaccine. Some people may have a rare but serious allergic reaction called anaphylaxis. If anaphylaxis happens, you will get medicine to treat the symptoms.

It is rare to have a serious side effect after a vaccine. Call Health Link at 811 to report any serious or unusual side effects.

Rare events after getting the

AstraZeneca/COVISHIELD vaccine

There have been very rare reports of blood clots, low levels of platelets (these help your blood to clot), and bleeding after getting the AstraZeneca/COVISHIELD vaccine. These events happened 4 to 28 days after getting the vaccine. The risk of these events after getting the AstraZeneca/COVISHIELD vaccine is about:

- 1 in 55,000 after the first dose
- 1 in 600,000 after the second dose

Research is still ongoing to find out more about the risk of these events after getting the AstraZeneca/COVISHIELD vaccine.

If you have any of the following symptoms within 42 days of being immunized, get medical help **right away:**

- trouble talking or moving a part of your body
- shortness of breath
- chest pain
- severe swelling, pain, or a colour change in your arm or leg
- stomach pain that does not go away
- a severe headache that does not go away
- blurry vision
- seizures

There have also been very rare reports of capillary leak syndrome (CLS) within the first few days after getting the AstraZeneca/COVISHIELD vaccine. In some reported cases, the person had a history of CLS. CLS is a serious condition that causes fluid to leak from the small blood vessels (capillaries). This causes sudden swelling of the arms and legs, sudden weight gain, and low blood pressure causing you to feel faint. CLS can be fatal. Get medical help right away if you have any of these symptoms.

Research is still ongoing to find out more about the risk of CLS after getting AstraZeneca/COVISHIELD vaccine.

Rare events after getting either the Pfizer-BioNTech or the Moderna vaccine

There have been very rare reports of myocarditis (inflammation of the heart muscle) and pericarditis (inflammation of the lining around the heart) within 7 days of getting either the Pfizer-BioNTech or the Moderna vaccine. This inflammation can cause shortness of breath, chest pain or pressure, or a very fast or abnormal heart rate. Get medical help right away if you have any of these symptoms. These rare events were reported mostly after the second dose and in young adults and adolescents. Most cases were mild and got better with treatment. The Moderna vaccine may have a higher risk of these events than the Pfizer-BioNTech vaccine. Because the Moderna vaccine was only recently approved for use in 12 to 17 year olds, there is more to learn about the risk of these rare events for this age group after getting that vaccine. Research is still happening to find out more about the risk of these events from mRNA COVID-19 vaccines. Talk to your healthcare provider, if you have ever had myocarditis or pericarditis and have questions about getting a COVID-19 vaccine. At this time, there is not enough information to know if there is a higher risk of these events in these groups.

Your risk of serious illness from COVID-19 is much higher than it is for developing a rare event after getting this vaccine.

Go to COVID-19 vaccine – Frequently Asked Questions on <u>ahs.ca/topics/Page17389.aspx</u> for more information.

How can I manage side effects?

- To help with soreness and swelling, put a cool, wet cloth over the area where you had the needle.
- There is medicine to help with a fever or pain. Check with your doctor or pharmacist if you are not sure what medicine or dose to take. Follow the directions on the package.
- Children under the age of 18 years should **not** take aspirin because it can cause serious health problems.
- Some people with health problems, such as a weak immune system, must call their doctor if they get a fever. If you have been told to do this, call your doctor even if you think the fever is from the vaccine.

What if I had or am getting another type of vaccine?

You can get an inactivated (non-live) vaccine at the same time or any time before or after getting a COVID-19 vaccine. If you have had a live vaccine, you need to wait 14 days before you get a COVID-19 vaccine. Wait at least 14 days after you have a dose of the COVID-19 vaccine before you get a live vaccine.

Who should not get a COVID-19 vaccine?

You may not be able to get a COVID-19 vaccine if you:

- have an allergy to any part of the vaccine
- had a severe (serious) or unusual side effect after this vaccine or one like it
- are under age 11 years
- have had a live vaccine in the last 2 weeks

Check with your doctor or a public health nurse before you get the vaccine. Be sure to talk to your doctor **before** you get a COVID-19 vaccine, if you:

- have a weak immune system (because of a medicine you take or a health problem)
- have an autoimmune disorder (like rheumatoid arthritis or lupus)
- have had a stem cell or organ transplant
- have a history of capillary leak syndrome (CLS)
- have a history of myocarditis or pericarditis after receiving a dose of COVID-19 vaccine

Always tell your healthcare provider if you have allergies or if you have had a side effect from a vaccine in the past.

For More Information

S Call Health Link at 811

Go to MyHealth.Alberta.ca G_0

Go to ImmunizeAlberta.ca Go to ahs.ca/covidvaccine