

For use at Alberta Health Services (AHS) immunization programs. Use this form when a parent or alternate decision-maker is **not** able to be with the person being immunized at an AHS immunization service. An alternate decision-maker could be an agent, guardian, specific decision-maker or co-decision-maker. The parent or alternate decision-maker should **complete this form** and **send it** with the person being immunized, to the AHS immunization service. Alternate decision-makers must also include a copy of the documents showing they are authorized as alternate decision-maker. These documents are required for this person to be immunized.

Personal information for the person being immunized		
Name (Last, First, Middle)		
Date of Birth (dd-Mon-yyyy)	Personal Health Number (PHN)	
Health information for the person being immunized (If you need more space, use the other side of this form.)		
Does this person have any allergies, including allergies to any vaccine, medicine, or food?		<input type="checkbox"/> No <input type="checkbox"/> Yes
If yes, describe _____		
Does this person have any chronic illness? (List all if more than one)		<input type="checkbox"/> No <input type="checkbox"/> Yes
If yes, describe _____		
Has this person ever had capillary leak syndrome?		<input type="checkbox"/> No <input type="checkbox"/> Yes
Has this person ever had immune thrombocytopenia?		<input type="checkbox"/> No <input type="checkbox"/> Yes
If this person is under age 18 years, have they ever had multisystem inflammatory syndrome (MIS-C)?		
<input type="checkbox"/> No <input type="checkbox"/> Yes <input type="checkbox"/> Not applicable		
Is this person taking any medicine?	<input type="checkbox"/> No <input type="checkbox"/> Yes	If yes, describe _____
Is this person pregnant?	<input type="checkbox"/> No <input type="checkbox"/> Yes	Is this person breastfeeding?
	<input type="checkbox"/> No <input type="checkbox"/> Yes	<input type="checkbox"/> No <input type="checkbox"/> Yes
Has this person had COVID-19 vaccine before?		<input type="checkbox"/> No <input type="checkbox"/> Yes If yes, when? _____
Has this person ever had a side effect from COVID-19 immunization?		<input type="checkbox"/> No <input type="checkbox"/> Yes
If yes, describe _____		
Has this person had myocarditis or pericarditis within 6 weeks of getting a COVID-19 vaccine?		<input type="checkbox"/> No <input type="checkbox"/> Yes
Has this person had a COVID-19 infection?		<input type="checkbox"/> No <input type="checkbox"/> Yes If yes, when? _____
Has this person had any vaccines in the last 4 weeks?		<input type="checkbox"/> No <input type="checkbox"/> Yes
If yes, list vaccine and date _____		
<p>I confirm that I have read the attached COVID-19 vaccine information. I know about and understand the risks, benefits, and common side effects of this vaccine. Any questions I may have had about this person getting this vaccine have been answered by calling the local public health office or Health Link at 811. I understand the information I have been given. I understand this consent is for all doses of the vaccine. I will contact the local public health office or the healthcare provider giving the COVID-19 vaccine if this person:</p> <ul style="list-style-type: none"> • has any changes to their health before getting any dose of the COVID-19 vaccine • has a severe or unusual side effect after any dose of the COVID-19 vaccine (other than the expected side effects listed on the COVID-19 vaccine information sheet) <p>I consent to this person getting the (only select one):</p> <p><input type="checkbox"/> Pfizer-BioNTech Comirnaty Original (Non-Bivalent) mRNA COVID-19 vaccine (primary series for ages 6 months and older)</p> <p><input type="checkbox"/> Moderna SpikeVax Original (Non-Bivalent) mRNA COVID-19 vaccine (primary series for ages 6 months to 5 years)</p> <p><input type="checkbox"/> Pfizer-BioNTech Comirnaty Omicron-Containing Bivalent mRNA COVID-19 vaccine (booster dose for ages 5 years and older)</p> <p><input type="checkbox"/> Either the Pfizer-BioNTech (Original or Bivalent) or Moderna (Original) mRNA COVID-19 vaccine (Depending which vaccine is recommended for this person at the time of their immunization)</p> <p><input type="checkbox"/> Novavax (Nuvaxovid) COVID-19 vaccine (for ages 12 years and older)</p> <p><input type="checkbox"/> Janssen (Johnson & Johnson) viral vector-based COVID-19 vaccine (for ages 18 years and older)</p> <p>I understand that I may withdraw this consent at any time by calling the local public health office or healthcare provider giving the COVID-19 vaccine.</p> <p>I confirm that I have the legal authority to consent to this immunization.</p>		
Printed name of person giving consent	Daytime phone	Other phone
<p>Relationship to person</p> <p><input type="checkbox"/> Parent (with legal authority to consent) <input type="checkbox"/> Guardian/Legal representative <input type="checkbox"/> Agent</p> <p><input type="checkbox"/> Co-decision-maker <input type="checkbox"/> Specific decision-maker <input type="checkbox"/> Other _____</p>		
Signature of person giving consent		Date (dd-Mon-yyyy)

Alberta Health Services collects health information according to Section 20 of the Health Information Act (HIA). This information is used to provide health services, determine eligibility for health services, or to carry out any other purpose authorized by the HIA. If you have any questions about this, please ask the healthcare provider giving the immunization or contact your local public health office. If you do not know your local public health office, call Health Link at 811 to get this information.



For Office Use Only

[illegible]

COVID-19 vaccines – mRNA (age 6 months and older)

What are COVID-19 vaccines?

COVID-19 vaccines protect against the SARS-CoV-2 virus (also known as COVID-19). The virus causes an infection in the lungs and airways and is a type of respiratory illness. In some cases, the infection can cause problems with other organs or other parts of the body. Go to ahs.ca/covid to learn more about COVID-19.

The following COVID-19 vaccines are approved for use in Canada:

- **mRNA vaccines:** Pfizer-BioNTech (Comirnaty Original [Non-Bivalent] and Comirnaty Bivalent) and Moderna (SpikeVax Original [Non-Bivalent] and SpikeVax Bivalent)
- **viral vector-based vaccines:** AstraZeneca (Vaxzevria)/COVISHIELD and Janssen (Johnson & Johnson)
- **protein-based vaccine:** Novavax (Nuvaxovid)
- **plant-based vaccine:** Medicago (Covifenz)

The Pfizer-BioNTech Original (Non-Bivalent) and Bivalent, Moderna Original (Non-Bivalent), Janssen, and Novavax vaccines are available in Alberta. For information about the protein-based and viral vector-based vaccines, read the vaccine information sheets on [ImmunizeAlberta.ca](https://immunizealberta.ca).

Who should get an mRNA COVID-19 vaccine?

Everyone is at risk of COVID-19. You should get an mRNA COVID-19 vaccine if you are age 6 months or older. COVID-19 vaccines are free.

Do I need a vaccine if I have already had COVID-19?

If you have already had COVID-19, it is not yet known how long your protection will last or how much protection you will have against variants. It is important to get a COVID-19 vaccine even if you have already had the virus.

The vaccine may give you better protection if you wait a while after having COVID-19 and then get a vaccine. How long to wait depends on your health history, the number of doses of COVID-19 vaccine you have had, and your risk of getting very sick from COVID-19.

If you have had COVID-19 in the past, check with your healthcare provider about when to get a COVID-19 vaccine.

What if my child is getting a COVID-19 vaccine?

Consent for a COVID-19 vaccine for children under age 18 years is provided by a parent or guardian. If a parent or guardian cannot be at the appointment, they can give consent in writing using the consent form at ahs.ca/VaccineUnder18. In some cases, children under age 18 years may be able to give their own consent.

How well do COVID-19 vaccines work?

How well the vaccines work against COVID-19 is different for each variant of the virus. COVID-19 vaccines are the best way to lower your risk of getting COVID-19 or getting very sick from it.

The mRNA vaccines give the best protection. The Pfizer-BioNTech Bivalent vaccine is the only vaccine available in Alberta that is made to protect against Omicron variants, which are the strains that are currently spreading. The protein-based vaccine gives more protection than the viral vector-based vaccines. However, all the vaccines work very well to lower your risk of getting seriously ill and needing to be in the hospital.

What vaccine can I get?

Primary series

The primary series of a vaccine is the initial number of doses you need to get protection. The vaccine used for your primary series depends on your age:

- If you are age 6 months to 5 years, you can get either the Pfizer-BioNTech or Moderna Original (Non-Bivalent) vaccines for your primary series.
- If you are age 6 years or older, you can get the Pfizer-BioNTech Original (Non-Bivalent) vaccine for your primary series.

If you are age 6 months to 4 years, it will take:

- 16 to 20 weeks to complete your primary series with the Pfizer-BioNTech vaccine, or
- 8 to 12 weeks to complete your primary series with the Moderna vaccine

The Moderna vaccine is recommended for children age 6 months to 4 years who have a weak immune system to give them protection as quickly as possible.

If you are age 5 years, it is recommended to get the Pfizer-BioNTech vaccine over the Moderna vaccine. The Pfizer-BioNTech vaccine has been shown to have a lower risk of myocarditis and pericarditis, especially in people age 12 to 29 years. It may also have a lower risk for children age 5 to 11 years.

If you cannot get or do not want an mRNA vaccine, you can get the protein-based vaccine if you are age 12 years or older or a viral vector-based vaccine if you are age 18 years or older.

Usually you get the same vaccine for all your doses. If your first-dose vaccine is not available or you cannot have it, you can get a different COVID-19 vaccine for your next dose. Whichever vaccine you get to complete your primary series protects you against COVID-19.

Booster doses

A booster dose will help to keep you protected for longer. You can get the Pfizer-BioNTech Bivalent vaccine for a booster dose if you are age 5 years and older. Children under age 5 years are not eligible for a booster dose at this time.

If you cannot get or do not want a bivalent mRNA vaccine for a booster dose, you may be able to get the protein-based vaccine if you are age 12 years or older.

Talk to your healthcare provider about which vaccine you can get and when to have your doses.

How many doses of mRNA COVID-19 vaccine do I need?

Primary series

The number of doses in your primary series depends on your age, health, and the type of vaccine you get.

If you get the Moderna vaccine, you need at least 2 doses. If you get the Pfizer-BioNTech vaccine, you need at least 2 doses if you are age 5 years or older and at least 3 doses if you are age 6 months to 4 years.

For some people, these doses may not give enough protection. You may need an extra dose to complete your primary series if you have a health problem that weakens your immune system. For example:

- You have had or will have an organ or stem cell transplant.
- You have kidney disease and need dialysis.
- You have acquired immunodeficiency syndrome (AIDS).
- You take certain medicines that weaken your immune system.

Studies have shown that an extra dose in the primary series may give better protection to adults with a weak immune system. In babies and children with a weak immune system, an extra dose may also give better protection, but research is still happening to learn more. If you get a 4-dose primary series of the Pfizer-BioNTech vaccine, this is considered "off-label use." This means the vaccine is not licensed (approved for use) in this situation but is supported by vaccine experts.

Booster doses age 5 to 11 years

If you are age 5 to 11 years, you can get 1 booster dose.

If you got the original (non-bivalent) mRNA vaccine as a booster dose, you can get a bivalent mRNA booster dose if you have certain health problems, such as:

- a weak immune system due to an illness or medicine
- kidney disease and need dialysis
- heart, lung, or liver problems
- diabetes
- a lot of extra weight
- had or will have an organ or stem cell transplant

Booster doses age 12 years and older

If you are age 12 years or older, you can get 1 bivalent mRNA booster dose on or after September 21, 2022.

If you are at high risk of severe illness from COVID-19, you can get a second bivalent mRNA booster dose. You may be at high risk if:

- You are age 65 years or older.
- You are age 18 years or older and live in a long-term care home or supportive living facility (such as a nursing home or seniors' lodge).
- You are age 18 years or older and have certain health problems such as a weak immune system due to an illness or medicine, you have had or will have an organ or stem cell transplant, or you have kidney disease and need dialysis.

When should I get my booster dose?

You need to wait 6 months after getting your last dose of COVID-19 vaccine or being infected with COVID-19 before getting a booster dose. Waiting 6 months helps the booster dose give you better protection.

You can get your booster dose 3 months after your last dose of COVID-19 vaccine or being infected with COVID-19 if:

- You live in a long-term care home or supportive living facility.
- You have had a stem cell transplant or are getting CAR T-cell therapy (a type of treatment for certain cancers).

If you get a second bivalent booster dose or you are age 5 to 11 years and get a Pfizer-BioNTech Bivalent booster dose sooner than 6 months after your last dose, this is considered “off-label use”. Vaccine experts support using these vaccines in this way.

Talk to your healthcare provider about when you should get your booster dose.

Go to alberta.ca/covid19-vaccine to book an appointment for your booster dose.

Can I get an mRNA COVID-19 vaccine if I am pregnant or breastfeeding?

Yes, you can get an mRNA COVID-19 vaccine if you are pregnant or breastfeeding. Research shows that mRNA vaccines are the safest type of COVID-19 vaccines to get during pregnancy, which includes the bivalent mRNA vaccines.

When you are pregnant, you have a higher risk of getting very sick from COVID-19. Getting a COVID-19 vaccine lowers your risk of getting seriously ill from the virus.

If you are pregnant or breastfeeding and have questions about getting a COVID-19 vaccine, talk to your healthcare provider.

Where can I get a COVID-19 vaccine?

Go to ahs.ca/covidvaccine to find out where and when you can get a COVID-19 vaccine. COVID-19 vaccines are free.

Are there side effects from mRNA COVID-19 vaccines?

There can be side effects from mRNA COVID-19 vaccines, but they tend to be mild and go away in a few days. Side effects may include:

- redness, swelling, or feeling sore where you had the needle
- feeling tired or have a headache
- a fever or chills
- body aches or sore joints
- pain in your arms or legs
- feeling sick to your stomach (nausea), vomiting (throwing up), or loose stool (diarrhea)
- swollen lymph nodes
- swelling or feeling sore in your armpit or groin
- a rash or hives

Children age 5 years and younger may also get upset easily or cry, have an earache, or may not want to eat.

At least 1 out of 100 people who got this vaccine reported 1 or more of these side effects. In some cases, it is unknown if the vaccine caused these side effects.

Current information shows that there is similar risk of side effects after each dose of an mRNA COVID-19 vaccine. Research continues to find out more about the risk of side effects after additional doses.

It is important to stay at the clinic for 15 minutes after your vaccine. Some people may have a rare but serious allergic reaction called anaphylaxis. If anaphylaxis happens, you will get medicine to treat the symptoms.

It is rare to have a serious side effect. Call Health Link at 811 to report any serious or unusual side effects.

What rare events have been reported after getting an mRNA COVID-19 vaccine?

There have been very rare reports of myocarditis (inflammation of the heart muscle) and pericarditis (inflammation of the lining around the heart) within 7 days of getting either the Pfizer-BioNTech or the Moderna vaccine. Most reported cases were mild and got better with treatment. The inflammation can cause shortness of breath, chest pain or pressure, or a very fast or abnormal heart rate. Get medical help **right away** if you have any of these symptoms.

Available information shows that these rare events were reported more commonly:

- after the second dose
- in those age 12 to 29 years
- in males
- in those who got the original (non-bivalent) Moderna vaccine

Research has shown that children age 5 to 11 years may also have a lower risk of these events after their primary series with Pfizer-BioNTech vaccine than adolescents and young adults.

In clinical trials for both the Pfizer-BioNTech and Moderna vaccines, there were no reports of myocarditis or pericarditis in children age 6 months to 5 years or in adults who got the bivalent vaccines.

Research is still happening to learn more about the risk of these events after a getting a bivalent vaccine and the risks in babies and children.

It is not known if having a history of myocarditis or pericarditis puts you at higher risk of having these rare events after a COVID-19 vaccine. Talk to your doctor before you get a dose of COVID-19 vaccine if:

- You had myocarditis or pericarditis within 6 weeks of getting a COVID-19 vaccine.
- You have a history of myocarditis or pericarditis and you have questions about getting a COVID-19 vaccine.

Research is happening to learn more about the risks of these rare events. Your risk of getting seriously ill from COVID-19 is much higher than your risk of having a rare event after these vaccines.

Go to ahs.ca/topics/Page17389.aspx for more information.

How can I manage side effects?

- To help with soreness and swelling, put a cool, wet cloth over the area where you had the needle.
- There is medicine to help with a fever or pain. Check with your doctor or pharmacist if you are not sure what medicine or dose to take. Follow the directions on the package.
- Some people with health problems, such as a weak immune system, must call their doctor if they get a fever. If you have been told to do this, call your doctor even if you think the fever is from the vaccine.

What vaccine is recommended for people with a weak immune system?

Research has shown mRNA vaccines provide the best protection for people with a weak immune system. Talk to your healthcare provider about which vaccine is best for you.

What if I had or am getting another type of vaccine?

You can get most vaccines at the same time as, any time before, or any time after an mRNA COVID-19 vaccine, including routine vaccines and the influenza vaccine. However, certain vaccines have a waiting period. If you had another vaccine in the last 4 weeks, check with your healthcare provider about when you can get the mRNA COVID-19 vaccine.

Who should not get an mRNA COVID-19 vaccine?

You may not be able to get an mRNA COVID-19 vaccine if:

- You have an allergy to any part of the vaccine.
- You had a severe (serious) or unusual side effect after this vaccine or one like it.
- You are under age 6 months.

If you have allergies or have had a side effect to this vaccine, check with your doctor or a public health nurse before you get the vaccine.

Although you can get the vaccine if you have a mild illness such as a cold or fever, you should stay home until you are feeling better to prevent spreading your illness to others.

Check with your healthcare provider about when you can get an mRNA COVID-19 vaccine if:

- You had COVID-19 in the past.
- You are under age 18 years and have a history of multisystem inflammatory syndrome (MIS-C).

Be sure to talk to your doctor **before** you get an mRNA COVID-19 vaccine if:

- You have a weak immune system (because of a medicine you take or a health problem).
- You have an autoimmune disorder (like rheumatoid arthritis or lupus).
- You have had a stem cell or organ transplant.
- You have a history of myocarditis or pericarditis within 6 weeks of getting a dose of COVID-19 vaccine.

COVID-19 vaccines - protein-based (age 12 years and older)

What are COVID-19 vaccines?

COVID-19 vaccines protect against the SARS-CoV-2 virus (also known as COVID-19). The virus causes an infection in the lungs and airways and is a type of respiratory illness. In some cases, the infection can cause problems with other organs or other parts of the body. Go to ahs.ca/covid to learn more about COVID-19.

The following COVID-19 vaccines are approved for use in Canada:

- **mRNA vaccines:** Pfizer-BioNTech (Comirnaty Original [Non-Bivalent] and Comirnaty Bivalent) and Moderna (SpikeVax Original [Non-Bivalent] and SpikeVax Bivalent)
- **viral vector-based vaccines:** AstraZeneca (Vaxzevria)/COVISHIELD and Janssen (Johnson & Johnson)
- **protein-based vaccine:** Novavax (Nuvaxovid)
- **plant-based vaccine:** Medicago (Covifenz)

The Pfizer-BioNTech Original (Non-Bivalent) and Bivalent, Moderna Original (Non-Bivalent), Janssen, and Novavax vaccines are available in Alberta. For information about the mRNA and viral vector-based vaccines, read the vaccine information sheets on [ImmunizeAlberta.ca](https://immunizealberta.ca).

Who can get the protein-based COVID-19 vaccine?

Everyone is at risk of COVID-19. You can get a protein-based COVID-19 vaccine if you are age 12 years or older and you cannot have or do not want an mRNA COVID-19 vaccine. COVID-19 vaccines are free.

Do I need a vaccine if I have already had COVID-19?

If you have already had COVID-19, it is not yet known how long your protection will last or how much protection you will have against variants. It is important to get a COVID-19 vaccine even if you have already had the virus.

The vaccine may give you better protection if you wait a while after having COVID-19 and then get a vaccine. How long to wait depends on your health history, the number of doses of COVID-19 vaccine you have had, and your risk of getting very sick from COVID-19.

If you have had COVID-19 in the past, check with your healthcare provider about when to get a COVID-19 vaccine.

How well do COVID-19 vaccines work?

How well the vaccines work against COVID-19 is different for each variant of the virus. COVID-19 vaccines are the best way to lower your risk of getting COVID-19 or getting very sick from it.

The mRNA vaccines give the best protection. The Pfizer-BioNTech Bivalent vaccine is the only vaccine available in Alberta that is made to protect against Omicron variants, which are the strains that are currently spreading. The protein-based vaccine gives more protection than the viral vector-based vaccines. However, all the vaccines work very well to lower your risk of getting seriously ill and needing to be in the hospital.

What vaccine can I get?

Primary series

The primary series of a vaccine is the initial number of doses you need to get protection.

It is recommended that you get an original (non-bivalent) mRNA COVID-19 vaccine for your primary series. If you cannot get or do not want an original (non-bivalent) mRNA vaccine, you can get the protein-based vaccine if you are age 12 years or older, or you can get the viral vector-based vaccine if you are age 18 years or older. Usually you get the same

vaccine for all your doses. If your first-dose vaccine is not available or you cannot have it, you can get a different COVID-19 vaccine for your next dose.

Whichever vaccine you get to complete your primary series protects you against COVID-19.

Booster doses

A booster dose helps to keep you protected for longer.

It is recommended that you get a bivalent mRNA vaccine for your booster dose. If you cannot get or do not want a bivalent mRNA vaccine, you may be able to get the protein-based vaccine if you are age 12 years or older.

Talk to your healthcare provider about which vaccine is best for you.

Go to ahs.ca/seconddose for more information.

How many doses of the protein-based COVID-19 vaccine do I need?

Primary series

You need at least 2 doses in your primary series of the protein-based COVID-19 vaccine.

For some people, 2 doses may not give enough protection. You may need a third dose to complete your primary series if you have a health problem that weakens your immune system. For example:

- You have had or will have an organ or stem cell transplant.
- You have kidney disease and need dialysis.
- You have acquired immunodeficiency syndrome (AIDS).
- You take certain medicines that weaken your immune system.

Studies have shown that a third dose may provide better protection for adults with a weak immune system. In children with a weak immune system, an extra dose may also give better protection, but research is still happening to learn more. If you get a 3-dose primary series of the protein-based vaccine, this is considered “off-label use”. This means the vaccine is not licensed (approved for use) in this situation but is supported by vaccine experts.

Booster doses

You can get up to 2 booster doses of the protein-based vaccine, but if you got a protein-based vaccine as a booster dose on or after September 21, 2022, then no further doses of the protein-based vaccine are recommended. However, you can still get at least 1 booster dose of a bivalent mRNA vaccine, or 2 doses if you are at high risk of severe illness.

When should I get my booster dose?

You need to wait 6 months after getting your last dose of COVID-19 vaccine or being infected with COVID-19 before you get a booster dose. Waiting 6 months helps the booster dose give you better protection. You can get your booster dose 3 months after your last dose of COVID-19 vaccine or being infected with COVID-19 if you have had a stem cell transplant or are getting CAR-T cell therapy (a type of treatment for certain cancers).

If you get a booster dose of the protein-based vaccine and it is sooner than 6 months after your last dose, or if you have already had a protein-based vaccine booster dose, or if you are age 12 to 17 years of age, it is considered “off-label use”. Vaccine experts support using the vaccine this way.

Talk to your healthcare provider about when you should get your booster dose. Go to alberta.ca/covid19-vaccine to book an appointment for your booster dose.

Can I get the protein-based COVID-19 vaccine if I am pregnant or breastfeeding?

Yes, you can get a protein-based COVID-19 vaccine if you are pregnant or breastfeeding. However, research shows that mRNA vaccines are the safest type of COVID-19 vaccines to get during pregnancy. There is more to learn about getting a protein-based COVID-19 vaccine when you are pregnant or breastfeeding.

When you are pregnant, you have a higher risk of getting very sick from COVID-19. Getting a COVID-19 vaccine lowers your risk of getting seriously ill from the virus.

If you are pregnant or breastfeeding, it is recommended that you talk to your healthcare provider before getting a protein-based COVID-19 vaccine.

Where can I get a COVID-19 vaccine?

Go to ahs.ca/covidvaccine to find out where and when you can get a COVID-19 vaccine. COVID-19 vaccines are free.

Are there side effects from the protein-based COVID-19 vaccine?

There can be side effects from the protein-based COVID-19 vaccine, but they tend to be mild and go away in a few days. Side effects may include:

- redness, swelling, or feeling sore where you had the needle
- headache, feeling tired or unwell
- a fever or chills
- body aches, sore joints, or pain in your legs or arms
- feeling sick to your stomach (nausea) or vomiting (throwing up)

At least 1 out of 100 people who got this vaccine reported 1 or more of these side effects. In some cases, it is unknown if the vaccine caused these side effects.

Current information shows that you are more likely to have these side effects after your second dose of the protein-based vaccine.

Research continues to find out more about the risk of side effects after additional doses.

It is important to stay at the clinic for 15 minutes after your vaccine. Some people may have a rare but serious allergic reaction called anaphylaxis. If anaphylaxis happens, you will get medicine to treat the symptoms.

It is rare to have a serious side effect. Call Health Link at 811 to report any serious or unusual side effects.

What rare events have been reported after getting the protein-based COVID-19 vaccine?

There have been very rare reports of myocarditis (inflammation of the heart muscle) and pericarditis (inflammation of the lining around the heart) shortly after getting the protein-based vaccine. The reported cases were mild and got better with treatment. It is still not known if these events were caused by the vaccine, and research is happening to learn more about the risk of these rare events after getting the protein-based vaccine.

The inflammation can cause shortness of breath, chest pain or pressure, or a very fast or abnormal heart rate. Get medical help **right away** if you have any of these symptoms.

It is not known if having a history of myocarditis or pericarditis puts you at higher risk of having these rare events after a COVID-19 vaccine. Talk to your doctor before you get a dose of COVID-19 vaccine if:

- You had myocarditis or pericarditis within 6 weeks of getting a COVID-19 vaccine.
- You have a history of myocarditis or pericarditis and you have questions about getting a COVID-19 vaccine.

Your risk of getting seriously ill from COVID-19 is much higher than your risk of having a rare event after these vaccines.

Go to COVID-19 vaccine – Frequently Asked Questions on ahs.ca/topics/Page17389.aspx for more information.

How can I manage side effects?

- To help with soreness and swelling, put a cool, wet cloth over the area where you had the needle.
- There is medicine to help with a fever or pain. Check with your doctor or pharmacist if you are not sure what medicine or dose to take. Follow the directions on the package.
- Some people with health problems, such as a weak immune system, must call their doctor if they get a fever. If you have been told to do this, call your doctor even if you think the fever is from the vaccine.

What vaccine is recommended for people with a weak immune system?

Research has shown mRNA vaccines provide the best protection for people with a weak immune system. Talk to your healthcare provider about which vaccine is best for you.

What if I had or am getting another type of vaccine?

You can get most vaccines at the same time as, any time before, or any time after the protein-based COVID-19 vaccine. However, certain vaccines have a waiting period. If you had another vaccine in the last 2 weeks, check with your healthcare provider about when you can get the protein-based COVID-19 vaccine.

Who should not get the protein-based COVID-19 vaccine?

You may not be able to get the protein-based COVID-19 vaccine if:

- You have an allergy to any part of the vaccine.
- You had a severe (serious) or unusual side effect after this vaccine or one like it.
- You are under age 12 years.

If you have allergies or have had a side effect to this vaccine, check with your doctor or a public health nurse before you get the vaccine.

Although you can get the vaccine if you have a mild illness such as a cold or fever, you should stay home until you are feeling better to prevent spreading your illness to others.

Check with your healthcare provider about when you can get the protein-based COVID-19 vaccine if you had COVID-19 in the past.

Be sure to talk to your doctor **before** you get the protein-based COVID-19 vaccine if:

- You have a weak immune system (because of a medicine you take or a health problem).
- You have an autoimmune disorder (like rheumatoid arthritis or lupus).
- You have had a stem cell or organ transplant.
- You have a history of myocarditis or pericarditis within 6 weeks of getting a dose of COVID-19 vaccine.

For more information about immunization



Call Health Link at 811



Scan the QR code or go to ImmunizeAlberta.ca



Scan the QR code or go to MyHealth.Alberta.ca

COVID-19 vaccines - viral vector-based (ages 18 years and older)

What are COVID-19 vaccines?

COVID-19 vaccines protect against the SARS-CoV-2 virus (also known as COVID-19). The virus causes an infection in the lungs and airways and is a type of respiratory illness. In some cases, the infection can cause problems with other organs or other parts of the body. Go to ahs.ca/covid to learn more about COVID-19.

The following COVID-19 vaccines are approved for use in Canada:

- **mRNA vaccines:** Pfizer-BioNTech (Comirnaty Original [Non-Bivalent] and Comirnaty Bivalent) and Moderna (SpikeVax Original [Non-Bivalent] and SpikeVax Bivalent)
- **viral vector-based vaccines:** AstraZeneca (Vaxzevria)/ COVISHIELD and Janssen (Johnson & Johnson)
- **protein-based vaccine:** Novavax (Nuvaxovid)
- **plant-based vaccine:** Medicago (Covifenz)

The Pfizer-BioNTech Original (Non-Bivalent) and Bivalent, Moderna Original (Non-Bivalent), Janssen, and Novavax vaccines are available in Alberta. For information about the mRNA and protein-based vaccines, read the vaccine information sheets on immunizealberta.ca.

Who can get a viral vector-based COVID-19 vaccine?

Everyone is at risk of COVID-19. You can get a viral vector-based COVID-19 vaccine if you are age 18 years or older and you cannot have or do not want an mRNA or protein-based COVID-19 vaccine. COVID-19 vaccines are free.

Do I need a vaccine if I have already had COVID-19?

If you have already had COVID-19, it is not yet known how long your protection will last or how much protection you will have against variants. It is important to get a COVID-19 vaccine even if you have already had the virus.

The vaccine may give you better protection if you wait a while after having COVID-19 and then get a vaccine. How long to wait depends on your health history, the number of doses of COVID-19 vaccine you have had, and your risk of getting very sick from COVID-19.

If you have had COVID-19 in the past, check with your healthcare provider about when to get a COVID-19 vaccine.

How well do COVID-19 vaccines work?

How well the vaccines work against COVID-19 is different for each variant of virus. COVID-19 vaccines are the best way to lower your risk of getting COVID-19 or getting very sick from it. The mRNA vaccines give the best protection. The Pfizer-BioNTech Bivalent vaccine is the only vaccine available in Alberta that is made to protect against Omicron variants, which are the strains that are currently spreading. The protein-based vaccine gives more protection than the viral vector-based vaccines. However, all the vaccines work very well to lower your risk of getting seriously ill and needing to be in the hospital.

What vaccine can I get?

Primary series

The primary series of a vaccine is the initial number of doses you need to get protection. It is recommended that you get an original (non-bivalent) mRNA COVID-19 vaccine for your primary series. If you cannot get or do not want an original (non-bivalent) mRNA vaccine, you can get the protein-based vaccine if you are age 12 years or older, or you can get the viral vector-based vaccine if you are age 18 years or older.

Usually you get the same vaccine for all your doses. If your first-dose vaccine is not available or you cannot have it, you can get a different COVID-19 vaccine for your next dose.

Whichever vaccine you get to complete your primary series protects you against COVID-19.

Booster doses

A booster dose will help keep you protected for longer.

It is recommended that you get a bivalent mRNA vaccine for your booster dose. If you cannot get or do not want a bivalent mRNA vaccine you may be able to get the Janssen vaccine for your booster dose if you are age 18 years or older.

Talk to your healthcare provider about which vaccine is best for you.

Go to ahs.ca/seconddose for more information.

How many doses of viral vector-based COVID-19 vaccine do I need?

Primary series

If you got the AstraZeneca/COVISHIELD vaccine (no longer available in Alberta), you need a second dose of a COVID-19 vaccine to complete your primary series.

If you get the Janssen vaccine, at this time you need only 1 dose to complete your primary series.

For some people, 2 doses may not give enough protection. You may need a third dose to complete your primary series if you have a health problem that weakens your immune system. For example:

- You have had or will have an organ or stem cell transplant.
- You have kidney disease and need dialysis.
- You have acquired immunodeficiency syndrome (AIDS).
- You take certain medicines that weaken your immune system.

Studies have shown that a third dose may provide better protection for adults with a weak immune system.

Booster doses

You can get 1 booster dose of the Janssen vaccine if you are age 18 years or older and you have had only 1 previous dose of Janssen vaccine (no other COVID-19 vaccines). You need to wait at least 2 months after completing your primary series or a COVID-19 infection before getting 1 booster dose of Janssen vaccine. No more doses of the Janssen vaccine are recommended after the 1 booster dose. However, you can still get at least 1 booster dose of a bivalent mRNA vaccine, or 2 doses if you are at high risk of severe illness. Talk to your healthcare provider about when you should get your booster dose. Go to alberta.ca/covid19-vaccine to book an appointment for your booster dose.

Can I get a viral vector-based COVID-19 vaccine if I am pregnant or breastfeeding?

Yes, you can get a viral vector-based COVID-19 vaccine if you are pregnant or breastfeeding. However, research shows that mRNA vaccines are the safest type of COVID-19 vaccines to get during pregnancy. There is more to learn about getting a viral vector-based COVID-19 vaccine when you are pregnant or breastfeeding.

When you are pregnant, you have a higher risk of getting very sick from COVID-19. Getting a COVID-19 vaccine lowers your risk of getting seriously ill from the virus.

If you are pregnant or breastfeeding, it is recommended that you talk to your healthcare provider before getting a viral vector-based COVID-19 vaccine.

Where can I get a COVID-19 vaccine?

Go to ahs.ca/covidvaccine to find out where and when you can get a COVID-19 vaccine. COVID-19 vaccines are free.

Are there side effects from the viral vector-based COVID-19 vaccine?

There can be side effects from the Janssen COVID-19 vaccine, but they tend to be mild and go away in a few days. Side effects may include:

- redness, swelling, or feeling sore where you had the needle
- feeling tired
- a headache
- a fever or chills
- body aches or sore joints
- feeling sick to your stomach (nausea)

At least 1 out of 100 people who got this vaccine reported 1 or more of these side effects. In some cases, it is unknown if the vaccine caused these side effects.

Current information shows that there is a similar risk of side effects after a first, second, or additional dose of a viral vector-based COVID-19 vaccine.

Research continues to find out more about the risk of side effects after additional doses.

It is important to stay at the clinic for 15 minutes after your vaccine. Some people may have a rare but serious allergic reaction called anaphylaxis. If anaphylaxis happens, you will get medicine to treat the symptoms.

It is rare to have a serious side effect. Call Health Link at 811 to report any serious or unusual side effects.

What rare events have been reported after getting the AstraZeneca/COVISHIELD or the Janssen vaccines?

Research is still happening to learn more about the risk of rare events after getting the AstraZeneca/COVISHIELD (no longer available in Alberta) or Janssen vaccines. It is not yet known if having a history of certain health problems puts you at a higher risk of rare events after having these vaccines. Your risk of getting seriously ill from COVID-19 is much higher than your risk of having a rare event after these vaccines.

Go to COVID-19 vaccine – Frequently Asked Questions on ahs.ca/topics/Page17389.aspx for more information.

Blood clots, low platelets, bleeding

There have been very rare reports of blood clots, low levels of platelets (these help your blood to clot), and bleeding after getting the AstraZeneca/COVISHIELD or the Janssen vaccines.

These events happened 4 to 28 days after getting the vaccine. The risk of these events after getting the AstraZeneca/COVISHIELD vaccine is about:

- 1 in 26,000 to 1 in 100,000 after the first dose
- 1 in 520,000 after the second dose

The risk of these events after getting the Janssen vaccine is about 1 in 300,000. Talk to your healthcare provider if you have a condition that puts you at risk for blood clots or if you have ever had immune thrombocytopenia (ITP). ITP is when your immune system attacks the blood cells you need for normal blood clotting and causes bleeding.

If you have any of the following symptoms within 42 days of being immunized, get medical help **right away**:

- trouble talking or moving a part of your body
- shortness of breath
- chest pain
- severe swelling, pain, or a colour change in your arm or leg
- stomach pain that does not go away
- a severe headache that does not go away
- blurry vision
- confusion
- seizures
- bruising or bleeding that is not normal for you

Capillary leak syndrome

There have been very rare reports of capillary leak syndrome (CLS) within the first few days after getting the AstraZeneca/COVISHIELD or Janssen vaccines. In some reported cases, the person had a history of CLS. CLS is a serious condition that causes fluid to leak from the small blood vessels (capillaries). This causes sudden swelling of the arms and legs, sudden weight gain, and low blood pressure causing you to feel faint. CLS can cause death. Talk to your healthcare provider if you have any history of CLS. Get medical help **right away** if you have any of these symptoms.

Guillain-Barré syndrome

There have been very rare reports of Guillain-Barré syndrome (GBS) up to 25 days after getting the AstraZeneca/COVISHIELD or Janssen vaccines. GBS is a serious condition that causes pain or numbness, muscle weakness, and in severe cases, paralysis. Most people fully recover from GBS but some may continue to have symptoms. GBS can cause death. Get medical help **right away** if you have any of these symptoms.

How can I manage side effects?

- To help with soreness and swelling, put a cool, wet cloth over the area where you had the needle.
- There is medicine to help with a fever or pain. Check with your doctor or pharmacist if you are not sure what medicine or dose to take. Follow the directions on the package.
- Some people with health problems, such as a weak immune system, must call their doctor if they get a fever. If you have been told to do this, call your doctor even if you think the fever is from the vaccine.

What vaccine is recommended for people with a weak immune system?

Research has shown mRNA vaccines provide the best protection for people with a weak immune system. Talk to your healthcare provider about which vaccine is best for you.

What if I had or am getting another type of vaccine?

You can get most vaccines at the same time as, any time before, or any time after a viral vector-based COVID-19 vaccine. However, certain vaccines have a waiting period. If you had another vaccine in the last 2 weeks, check with your healthcare provider about when you can get the viral vector-based COVID-19 vaccine.

Who should not get a viral vector-based COVID-19 vaccine?

You may not be able to get a viral vector-based COVID-19 vaccine if:

- You have an allergy to any part of the vaccine.
- You had a severe (serious) or unusual side effect after this vaccine or one like it.
- You have a history of capillary leak syndrome (CLS).
- You are under age 18 years.

If you have allergies or have had a side effect to this vaccine, check with your doctor or a public health nurse before you get the vaccine.

Although you can get the vaccine if you have a mild illness such as a cold or fever, you should stay home until you are feeling better to prevent spreading your illness to others.

Check with your healthcare provider about when you can get a viral vector-based COVID-19 vaccine if you had COVID-19 in the past.

Be sure to talk to your doctor **before** you get a viral vector-based COVID-19 vaccine if:

- You have a weak immune system (because of a medicine you take or a health problem).
- You have an autoimmune disorder (like rheumatoid arthritis or lupus).
- You have had a stem cell or organ transplant.
- You have a history of immune thrombocytopenia (ITP).
- You are at higher risk of blood clots.

For more information about immunization



Call Health Link at **811**



Scan the QR code or go to
ImmunizeAlberta.ca



Scan the QR code or go to
MyHealth.Alberta.ca