

Guide for Outbreak Prevention and Control in Long Term Care, Designated Supportive Living and Hospice Sites

Includes Respiratory & Gastrointestinal Illness

© 2023 Alberta Health Services, Communicable Disease Control, Notifiable Disease and Outbreak Program



This copyright work is licensed under the [Creative Commons - Attribution-NonCommercial-NoDerivatives 4.0 International](https://creativecommons.org/licenses/by-nc-nd/4.0/). To view a copy of this license, see [Creative Commons — Attribution-NonCommercial-NoDerivatives 4.0 International — CC BY-NC-ND 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/). You are free to copy and distribute the work including in other media and formats for non-commercial purposes, as long as you attribute the work to Alberta Health Services, do not adapt the work, and abide by the other license terms. The license does not apply to AHS trademarks, logos or content for which Alberta Health services is not the copyright owner.

This material is intended for general information only and is provided on an “as is”, “where is” basis. Although reasonable efforts were made to confirm the accuracy of the information, Alberta Health Services does not make any representation or warranty, express, implied or statutory, as to the accuracy, reliability, completeness, applicability or fitness for a particular purpose of such information. This material is not a substitute for the advice of a qualified health professional. Alberta health Services expressly disclaims all liability for the use of these material, and for any claims, actions, demands or suits arising from such use.

INTRODUCTION	6
Land Acknowledgement	7
How to Use the Guide	7
DEFINITION OF TERMS & GLOSSARY	10
HOW TO CONTACT THE AHS PUBLIC HEALTH OUTBREAK TEAM TO REPORT ILLNESS	
15	
Initial point of contact for the Public Health Outbreak Team reporting	15
Contacting the Public Health Outbreak Team after site has reported initial cases or an outbreak is declared	15
SECTION 1: PREPARING FOR POTENTIAL OUTBREAKS	16
Zone Outbreak Preparation Resources	16
Roles and Responsibilities for Outbreak Preparation	16
SECTION 2: ASSESSING A POTENTIAL OUTBREAK.....	19
2.1 Surveillance and Threshold for Investigation	19
Table A: Determining when to report respiratory illness symptoms to CEIR	19
Table B: Determining when to report GI illness to CEIR	20
Stool Specimen Collection.....	20
Scenarios which do not meet the Respiratory Surveillance Reporting Definition	20
Specimen collection prior to meeting the respiratory surveillance reporting definition.....	21
Resource - Symptoms? What Do I Do to Prevent a Potential Outbreak?.....	21
2.2 Reporting once the respiratory surveillance or GI illness reporting threshold is met	21
Figure 1: Potential Outbreak Notification Algorithm*	22
Table C: Initial Data Required by the Public Health Outbreak Team.....	23
2.3 Reporting Additional or Subsequent Cases to Public Health Outbreak Team	23
Roles and Responsibilities: Assessing for and Reporting Potential Outbreaks	24
Attachment 2.1: Surveillance Case Tracking Sheet.....	26
SECTION 3: MANAGING A POTENTIAL OUTBREAK.....	27
3.1 Infection Prevention and Control Measures	27
Figure 2: Routine Practices and Additional Precautions	28
3.2 Administrative Measures	29
3.3 Resident Restrictions	29
3.4 Restrictions on Affected Unit/Site	30
3.5 Admissions/Transfers from Acute Care Site to an LTC, DSL or Hospice Site	30
3.6 Transfers from an Outbreak LTC, DSL or Hospice Site to an Acute Care Site.....	30
3.7 Group/Social Activities and Other Events	30
3.8 Nourishment Areas/Sharing of Food.....	30
3.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3).....	30
3.10 Visitors and Designated Family/Support Person(s) (DFSPs)	30
3.11 Volunteers	31
3.12 HCW/Staff Outbreak Control Measures (Including volunteers, students, and physicians)	31
Table D: Health Care Worker/Staff Testing and Work Restrictions	32
3.13 Specimen Collection.....	33
3.14 Enhanced Environmental Cleaning and Disinfection	33

Roles and Responsibilities for Managing a Potential Outbreak.....	34
Attachment 3.1: ProvLab Specimen Collection Guidance.....	37
SECTION 4: OUTBREAK DEFINITIONS & DETERMINING OUTBREAK TYPE	40
Table E: COVID-19 Illness	40
Table F: Influenza-Like-Illness (ILI)	40
Table G: Influenza Illness	41
Table H: Gastrointestinal Illness	41
SECTION 5: GENERAL RECOMMENDATIONS FOR CONFIRMED OUTBREAKS.....	42
5.1 Infection Prevention and Control Measures	42
5.2 Administrative Measures	42
5.3 Resident Restrictions	43
5.4 Restrictions to Admissions / Transfers / Discharges on Affected Units/Site	44
5.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site.....	44
5.6 Transfers from an Outbreak LTC, DSL or Hospice site to an Acute Care Site	45
5.7 Group/Social Activities and Non-Resident Events	45
5.8 Nourishment Areas/Sharing of Food.....	45
5.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3).....	46
5.10 Visitors and Designated Family/Support Person(s).....	46
5.11 Volunteers	46
5.12 HCW/Staff Outbreak Measures (including volunteers, students, physicians)	46
5.13 Specimen Collection.....	47
5.14 Enhanced Environmental Cleaning and Disinfection	47
Roles and Responsibilities for Confirmed Outbreaks.....	47
Attachment 5.1: Outbreak Signage.....	51
Attachment 5.2: Facility CDC Outbreak Daily Report Portal (RedCap)	52
SECTION 6: CONFIRMED COVID-19 OUTBREAK	53
COVID-19 Outbreak Definition	53
Duration of Outbreak	53
6.1 Infection Prevention and Control Measures	53
6.2 Administrative Measures	53
6.3 Resident Restrictions	54
6.4 Restrictions on Affected Unit/Site	54
6.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site.....	54
6.6 Transfers from an outbreak facility to an acute care site	54
6.7 Group / Social Activities and Other Events	55
6.8 Nourishment Areas / Sharing of Food.....	55
6.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3).....	55
6.10 Visitors and Designated Family/Support Person(s).....	55
6.11 Volunteers	55
6.12 HCW/Staff Outbreak Measures (Including volunteers, students, and physicians)	56
6.13 Specimen Collection.....	56
6.14 Enhanced Environmental Cleaning and Disinfection	56
Roles and Responsibilities for Confirmed Outbreaks.....	56
SECTION 7: CONFIRMED INFLUENZA-LIKE ILLNESS OUTBREAK	58
7A. Influenza-Like Illness (ILI) Outbreak Definition	58
7B. Syndromic or Pathogen Specific Influenza-Like Illness (ILI) Outbreaks.....	58

7C. Duration of Outbreak	58
Table I: Organisms Commonly Associated with Respiratory Illness.....	59
7.1 Infection Prevention and Control	62
7.2 Administrative Measures	62
7.3 Resident Restrictions	62
7.4 Restrictions to Admissions / Transfers / Discharges on Affected Unit / Site	62
7.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site.....	62
7.6 Transfers from an Outbreak Facility to an Acute Care Site	63
7.7 Group / Social Activities and Other Events	63
7.8 Nourishment Areas / Sharing of Food.....	63
7.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3).....	63
7.10 Visitors and Designated Support Person(s).....	63
7.11 Volunteers	63
7.12 HCW/Staff Outbreak Measures (Including volunteers, students, physicians).....	63
7.13 Specimen Collection.....	63
7.14 Enhanced Environmental Cleaning and Disinfection	63
Roles and Responsibilities for Confirmed Outbreaks	64
SECTION 8: CONFIRMED INFLUENZA OUTBREAK	65
Influenza Outbreak Definition	65
Duration of Outbreak	65
8.1 Infection Prevention and Control	65
8.2 Administrative Measures	65
8.3 Resident Restrictions	65
8.4 Restrictions on Affected Units/Site	66
8.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site.....	66
8.6 Transfers from an Outbreak Facility to an Acute Care Site	67
8.7 Group / Social Activities and Other Events	67
8.8 Nourishment Areas / Sharing of Food.....	67
8.9 Operating Adult Day Programs during an Outbreak.....	67
8.10 Visitors and Designated Support Person(s).....	67
8.11 Volunteers	67
8.12 HCW/Staff Outbreak Measures (Including volunteers, students, physicians).....	67
8.13 Specimen Collection.....	69
8.14 Enhanced Environmental Cleaning and Disinfection	69
Roles and Responsibilities for Confirmed Outbreaks.....	69
Attachment 8.1: MOH Notice to all Unimmunized HCW	71
Attachment 8.2 MOH Notice to HCW Immunized less than 14 days prior to declaration of Outbreak 72	
Attachment 8.3: Influenza Vaccine Decision Tool for HCW/staff.....	73
SECTION 9: CONFIRMED GASTROINTESTINAL ILLNESS OUTBREAK.....	74
GI Illness Case Definition	74
GI Illness Outbreak Definition.....	74
9.1 Infection Prevention and Control Measures	75
Figure 3: Additional Precautions for GI Illness	76
9.2 Administrative Measures	76
9.3 Resident Restrictions	76
9.4 Restrictions on Affected Units/Site	77
9.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site.....	77

9.6 Transfers from an Outbreak Facility to Acute Care	78
9.7 Group/Social Activities and other Events	78
9.8 Nourishment Areas / Sharing of Food.....	78
9.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3).....	79
9.10 Visitor and Designated Family/Support Person(s) (DFSPs).....	79
9.11 Volunteer Restrictions	79
9.12 HCW/Staff Outbreak Measures (Including volunteers, students, physicians).....	80
9.13 Specimen Collection.....	80
9.14 Enhanced Environmental Cleaning and Disinfection	80
9.15 Management of “Relapse” Cases	82
Attachment 9.1: Data Collection Tool for Gastrointestinal Illness Outbreak Management....	83
SECTION 10: CLOSING AN OUTBREAK.....	84
APPENDIX A: SAMPLE LETTERS AND FORMS FOR OUTBREAK SEASON	
PREPARATION.....	85
Sample letters	85
Sample Forms.....	85
Antiviral (Oseltamivir) Dosing Recommendations.....	85
Advance Prescriptions for Antiviral Medication during Influenza Outbreaks.....	87
Antiviral (Oseltamivir) Dosing Recommendations.....	89
Provision of Antiviral Medication During Influenza Outbreaks.....	90
Outbreak Preparation Letter	92
Important Notice to Staff about Influenza and COVID-19	95
Resident or Alternate Decision Maker Fall Letter.....	96
APPENDIX B: SYMPTOMS? WHAT DO I DO TO PREVENT A POTENTIAL OUTBREAK	97
APPENDIX C: OUTBREAK CHECKLISTS	98

Introduction

This guide is intended as an operational guide to practice. This document provides current best-practice and evidence-based guidance for outbreak control and management of respiratory and gastrointestinal (GI) illness in Long Term Care (LTC), Designated Supportive Living (DSL) and Hospice sites. Information within this document contains evidence-based best practice as outlined in the [Alberta Health \(AH\) Public Health Disease Management Guidelines](#) and in Public Health Agency of Canada documents. Content was written in consultation and close collaboration with the following partners:

- Alberta Health Services (AHS):
 - Infection Prevention & Control (IPC)
 - Workplace Health & Safety (WHS)
 - Seniors Health
- Covenant Health:
 - Infection Prevention & Control
 - Occupational Health & Safety

Congregate living settings are high-risk environments for communicable disease outbreaks due to communal living arrangements and close proximity between residents. The potential severity of an outbreak can be increased in these settings due to potential underlying factors and health conditions that could contribute to vulnerability of severe forms of infections in these settings.

This document discusses the roles of the Public Health Outbreak Team (which includes Medical Officers of Health, and Alberta Health Services (AHS) Zone Communicable Disease Control/Environmental Public Health Outbreak Teams); Infection Prevention and Control/Designate; Facility Administration/Facility Management or their Designate; Occupational Health/Workplace Health and Safety or their Designate; Onsite Staff; and the Provincial Laboratory for Public Health (ProvLab).

The roles and responsibilities of AH; AHS Zone Operations (Executive Director for Seniors Health, Area Managers / Directors, Directors for LTC, DSL and Hospice) AHS Provincial Seniors Health and Continuing Care are not discussed in this guide. If individuals or facilities require more information about the roles and responsibilities of these additional partners, they are recommended to seek information directly from the respective groups.

Infectious disease outbreaks occur year-round and in different settings, including LTC, DSL and Hospice sites. Outbreak management measures for respiratory and gastrointestinal illnesses outlined in this guide reflect a usual seasonal respiratory illness impact at the population level. Additional measures may be recommended where heightened levels of infection or newly emerging pathogens warrant increased surge levels of outbreak preparation, prevention, and management and are outside the scope of this guide.

Effective outbreak management requires a multidisciplinary approach and involves individuals with different responsibilities. In compliance with the [Alberta Continuing Care Health Services Standards \(Standard 11\)](#), Alberta Health Services (AHS) facilities and contracted service providers will develop and implement written procedures for identifying, reporting and investigating notifiable diseases, and controlling any suspect outbreaks within LTC, DSL and Hospice sites.

The notification of outbreaks and other infectious disease threats in Alberta is mandated under Section 26 of the provincial [Public Health Act](#), and each Medical Officer of Health (MOH) is accountable for outbreak investigation and management. Early recognition of unusual clusters of illness and swift actions in response to these episodes are essential for effective management of outbreaks. Any health practitioner or person in charge of an institution (Facility Operator/Designate) must notify the Public Health Outbreak team if there is a confirmed or suspected communicable disease outbreak or any other illness or health condition occurring at an unusually high rate.

Note: Only the minimum outbreak measures for viral respiratory or GI illnesses are outlined here. These standard recommendations may be enhanced or modified depending on the specific scenario.

If considering implementation of outbreak control measures beyond those recommended in this guide, it is important to consider the potential impact on the well-being of residents (e.g., if a resident is receiving end of life care, a conversation with the family about visitation during an outbreak may be needed). Exceptions are recommended in consultation with the Public Health Outbreak Team. For more information, consult the Public Health Outbreak Team.

In the event of an outbreak, or threat of an outbreak of an unusual infectious disease, (such as a new pandemic or any other infrequent infectious disease), direction on best practices for outbreak management will be provided by the MOH and may broaden the recommendations in this guide.

In instances of unusual clusters or high severity of a circulating disease in the Alberta population, additional population-wide measures may be recommended or required that exceed the standard outbreak management and prevention strategies outlined in this guide.

While it is recognized that *Clostridioides difficile* and multi-drug resistant organisms (e.g., MRSA, VRE), can be responsible for clusters or outbreaks, and that some of the measures outlined in this guide may be applicable in preventing or controlling them, it is beyond the scope of this document to include these organisms, due to their unique epidemiological properties.

Land Acknowledgement

We would like to recognize that our work takes place on historical and contemporary Indigenous lands, including the territories of Treaties 6, 7 & 8, and the homeland of the Métis. We also acknowledge the many Indigenous communities that have been forged in urban centres across Alberta.

How to Use the Guide

- Users are encouraged to familiarize themselves with the layout of the document. It is a comprehensive document that has been divided into eleven sections for user ease.
- Roles and responsibilities are presented for those that have a role in outbreak management. A user can refer to the information specific to their role and may review information about additional roles on an as-needed basis.
- To assist with using the document, the user can refer to the section that is appropriate to the type or stage of the outbreak. A brief description of each section is presented below.

- Due to the complex nature of LTC/DSL/Hospice settings, the individual fulfilling the roles and responsibilities within a site may vary. Refer to site roles and responsibilities.

[Section 1 Preparing for a Potential Outbreak](#)

- This section is to be used for review prior to the start of the outbreak season.
- [Appendix A](#) contains examples of the resources that the Public Health Outbreak Team will send out prior to the beginning of the outbreak season.
- Includes an overview of the roles and responsibilities for outbreak preparation and prevention.

[Section 2 Assessing a Potential Outbreak](#)

- This section is provided to help the user determine if a resident or staff member has any of the symptoms that would require reporting to the Public Health Outbreak Team via CEIR. It also includes how to contact CEIR.
- Includes an overview of the roles and responsibilities for assessing and reporting a potential outbreak.

[Section 3 Managing a Potential Outbreak](#)

- This section outlines the steps that need to be taken to contain the potential outbreak.
- Includes information on Infection Prevention and Control, Resident Restrictions, Visitors, Volunteers, Specimen Collection, etc.
- Includes an overview of the roles and responsibilities for managing the potential outbreak.

[Section 4 Outbreak Definition and Determining Outbreak Type](#)

- Includes case and outbreak definitions for COVID-19 Illness, Influenza-like Illness, Influenza Illness and GI Illness.

[Section 5 General Recommendations for Confirmed Outbreaks](#)

- This section provides general outbreak control strategies that are to be implemented regardless of the identified pathogen. These strategies are used in combination with [Section 3](#).
- Includes an overview of the roles and responsibilities for confirmed outbreaks.

[Section 6 Confirmed COVID-19 Outbreak](#)

- This section provides guidance for COVID-19-specific outbreak measures in addition to outbreak measures outlined in [Sections 3](#) and [5](#)
- Includes an overview of the roles and responsibilities for a confirmed COVID-19 outbreak.

[Section 7 Confirmed Influenza-like Illness \(ILI\) Outbreak](#)

- This section provides guidance for when no pathogen is identified or when a non-COVID-19, non-Influenza pathogen is identified.
- The outbreak control strategies in this section are implemented in addition to those in [Sections 3](#) and [5](#).

Section 8 Confirmed Influenza Outbreak

- This section provides guidance on managing an Influenza outbreak.
- The strategies in this section would be implemented in addition to those in [Sections 3](#) and [5](#).
- Provides direction on management of the unimmunized health care worker/staff.
- Provides information for the use of antivirals in an influenza outbreak.
- Includes an overview of the roles and responsibilities for a confirmed Influenza outbreak.

Section 9 Confirmed Gastrointestinal Illness Outbreak

- This section provides additional outbreak management measures when a GI Illness outbreak is declared.
- The strategies outlined in this section would be implemented in addition to those in [Sections 3](#) and [5](#)

Section 10 Closing an Outbreak

- This section provides direction on closing all types of outbreaks.

Definition of Terms & Glossary

Acute Care - Includes all urban and rural hospitals, psychiatric facilities, and urgent care facilities where inpatient care is provided.

Admission and transfer status - Determined in consultation with the Outbreak Management Team (OMT) and categorized as follows:

- **“Open”** - The facility/unit remains open to all resident admissions, transfers, and discharges.
- **“Restricted”** - Depending upon the circumstances and the infectious agent involved, admission and transfer status may range from NO admission to selected resident admissions, transfers and discharges as permitted under the direction of the zone Medical Officer of Health and in consultation with the OMT. This approach is intended to be flexible allowing for individual assessments to be made based on established criteria without undue risk to residents/program/system.

Adult Day Program (ADP) - Day program designed for adults over the age of 18 who may have physical and/or memory challenges or are living with a chronic illness. They play a key role in allowing people to remain living in the community as long as possible by optimizing their physical, spiritual, social, and emotional function. Adult day programs also provide respite and education for caregivers (e.g., CHOICE, C3, and Bridges).

AHS Home Care - Publicly-funded health care and support services provided to eligible residents as governed by the Alberta Home Care Program Regulations of the *Public Health Act*. These services are provided to individuals living with frailty, disability, acute or chronic illness living at home or in a supportive living setting as defined at AHS [Home Care](#).

AHS Home Care Operations - This term applies to the AHS Zone Seniors Health and Continuing Care staff that have responsibility to participate in outbreak management leadership at LTC, DSL and Hospice sites where home care services are provided. Examples might include:

- AHS Case Manager (CM) - a regulated health care provider who coordinates and integrates health care services within and across the health care system; facilitates access to and continuity of health services across the continuum of care (e.g., when a resident is hospitalized or discharged from one continuing care service/program to another; and assists residents in accessing appropriate services within the community.
- AHS Allied Health
- Transition Services/Coordinated Access - coordination of resident care and movement between sites and to different levels of care including but not limited to assessment for Home Care services, and assessment and admission to designated living options (LTC, DSL).

AHS Provincial Seniors Health and Continuing Care - AHS Provincial Seniors Health and Continuing Care focuses on improving health care services and practices to enable seniors to optimize their health, well-being, and independence. The provincial program works closely with stakeholders across the province to align and optimize services to meet the healthcare needs of Alberta's seniors based on evidence and best practices.

AHS Zone Operations - This term applies to the AHS Seniors Health staff that have responsibility to participate in outbreak management leadership at AHS owned and contracted sites.

Appropriate Mask - The type of mask (surgical, KN95, N95) recommended per the AHS IPC [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)

CEIR - AHS Coordinated Early Identification and Response line (CEIR): 1-844-343-0971. A provincial, centralized outbreak reporting and response line. This is a collaborative effort to assist with initial site support for facilities with vulnerable populations. CEIR assists with the implementation of initial outbreak management and control measures including initial specimen collection recommendations, isolation protocols, staffing, personal protective equipment (PPE), education, etc.

Close Contact - Any person suspected to have been exposed to an infected person or a contaminated environment to a sufficient degree to have had the opportunity to become infected or colonized with an organism.

Cluster - Aggregation of similar, relatively uncommon events or diseases in space and/or time in amounts that are believed or perceived to be greater than could be expected by chance.

Cohorting - Controlling the movement of HCW/staff and residents for the purpose of limiting an outbreak to a specific unit/floor/area within a larger facility (instead of having the outbreak declared facility wide). The Public Health Outbreak Team will assess specific circumstances to assist in determining the appropriate scope for the outbreak investigation.

Congregate setting - Refers to settings where residents receive care and/or services in a communal environment with other residents. Examples include Acute care, long term care, designated supportive living, hospice, adult day programs, non-designated supportive living sites including lodges and private retirement residence.

COVID-19 – For COVID-19 illness case and outbreak definitions refer to [Table E](#).

Contact and Droplet Precautions - see [Figure 2](#).

Designated Supportive Living (DSL) - A home-like setting where people can maintain control over their lives while also receiving the support they need. The buildings are specifically designed with common areas and features to allow individuals to “age in place”. Building features include private space and a safe, secure, and barrier-free environment. DSL promotes resident independence and aging in place through the provision of services such as 24-hour monitoring, emergency response, security, meals, housekeeping, and life-enrichment activities. Publicly funded personal care and health services are provided to supportive living residents based on assessed unmet needs.

- **Designated Supportive Living Level 3 (DSL3)** - Setting that provides accommodation, meals, housekeeping, linen, and recreational services where healthcare services are provided on a scheduled basis but can be accessed as needed. Twenty-four hour on-site scheduled and unscheduled personal care and support services are provided by Health Care Aides. Professional health services including Registered Nurse services with 24 hour on-call availability, case management and other consultative services are provided through AHS.

- **Designated Supportive Living Level 4 (DSL4)** - Living option that provides accommodation, meals, housekeeping, linen, and recreational services where a higher level of personal care supports, and health care services are provided onsite for scheduled and unscheduled care needs according to the individuals plan of care. Twenty-four hour on-site scheduled and unscheduled professional and personal care, and support services are provided by Licensed Practical Nurses and Health Care Aides. Professional health services including Registered Nurse services with 24 hour on-call availability, case management and other consultative services are provided through AHS.
- **Designated Supportive Living Level 4 Dementia (DSL4D)** - DSL4 that provides specialized dementia care.

Designated family/support person (DFSPs) - one or more individuals identified by the resident/patient as an essential support, and who the resident/patient wishes to be included in any encounters with the health care system, including, but not limited to, family, relatives, friends, and informal or hired caregivers. Refer to [Visiting a Loved One at an AHS Facility](#).

Drug Identification Number (DIN) - A number assigned by Health Canada to a drug product prior to being marketed in Canada.

Emergency Medical Services (EMS) Community Paramedic (CP) Program - A Mobile Integrated Healthcare Program designed to provide immediate or scheduled medical supervision and treatments that are currently unavailable in the community setting. This program works in collaboration with the resident's existing healthcare services. The target population are the medically fragile individuals requiring specialized treatments that are necessary to stay out of hospital, or support "at risk" individuals for readmission to hospital & /or visit to emergency (such as medical management during Outbreaks, which may include swab collection in some circumstances). A referral is required to access this service at the Zone level.

Exclusion - A measure that prevents symptomatic/infected/susceptible HCW/staff from working, until such time that the risk for residents or HCW/staff is low or minimal, as recommended by the Public Health Outbreak Team or Workplace Health and Safety or designate.

Exposure Investigation Number (EI Number) - A number assigned by the Provincial Laboratory for Public Health for the purpose of tracking laboratory specimens associated to a specific event (e.g., a potential outbreak) at a specific location and time.

Facility Operator/Facility Management - A formal leader within the physical site, responsible for the day-to-day operations of the site. This may or may not be an AHS employee depending on the type of facility.

Family-style meal service - Involves filling a common vessel, such as a tray or bowl, with a large portion of food and setting the vessel on the table allowing residents to serve themselves from the common vessel.

Gastrointestinal (GI) Illness - For GI illness case and outbreak definitions refer to [Table B](#).

Health Care Workers (HCW) - As defined by Alberta Health (AH) includes all health practitioners and all individuals (including nutrition and food services, housekeeping, recreation etc.) at increased risk for exposure to, and/or transmission of, a communicable disease because they work, study, or volunteer in one or more of the following health care environments: hospital, nursing home (facility living), supportive living accommodations, or home care setting, mental health facility, community setting, office or clinic of a health practitioner, clinical laboratory.

Health Care Workers/Staff (HCW/staff) - For the purpose of this document, the term HCW/staff will be used consistently, and the facility is responsible for determining if an individual is considered to be a HCW or a staff member. The facility will determine when actions need to be taken by all members at the site or only those that are considered to be a HCW.

- **Onsite Staff** – The term onsite staff is used for HCW/staff that in some capacity of their job work directly at that site. When an individual is on-site, they are expected to fulfill the Roles and Responsibilities of onsite HCW/staff as outlined in each section. The examples listed below are not an all-inclusive list:
 - Staff employed directly by the site
 - Staff who are contracted service providers and work at the site (e.g., Contracted Pharmacy Services)
 - AHS staff who work and provide services at the site.

Hospice - Any facility in which residential hospice services are offered or provided by Alberta Health Services or by a service provider under contract with Alberta Health Services.

Infection Control Designate (ICD) - Someone assigned to be accountable for IPC issues in a facility.

Infection Control Professional (ICP) - A health professional with specialized knowledge responsible for infection prevention and control within the facility or area of practice. ICPs come from several disciplines, including nursing, medicine, microbiology, medical technology and/or epidemiology and may be certified or working toward certification in infection control (CIC®).

Influenza-like Illness (ILI) - for ILI illness case and outbreak definitions refer to [Table F](#).

Influenza Illness – for influenza illness case and outbreak definitions refer to [Table G](#).

IPC - Infection Prevention and Control

Long Term Care (LTC) Facility - Long term care refers to a continuum of medical and social services designed to support the needs of people living with chronic health problems who require the oversight of a registered nurse 24-hours a day. Long term care services include traditional medical services, social services, and housing. Clients admitted into long-term care are required to pay accommodation fees (room and board and other costs associated) as set by government. Long term care may be referred to as auxiliary hospitals and nursing homes.

Most Responsible Health Practitioner - The **health practitioner** who has responsibility and accountability for the specific treatment/procedure(s) provided to a resident and who is authorized by Alberta Health Services and/or the facility to perform the duties required to fulfill the delivery of such a treatment/procedure(s) within the scope of his/her practice.

Occupational/Workplace Health & Safety (OHS/WHS) - Designated personnel responsible for staff health and safety in facilities. In some facilities, Employee Health or the Site Management or the Site Medical Leader may fill this role.

Outbreak - “The occurrence of cases of disease in excess of what would normally be expected in a defined community, geographical area or season” (World Health Organization, 2018). A common source of infection or the identification of transmission between cases are not required. The epidemiologic features of an outbreak and subsequent public health actions are assessed through the outbreak investigation process.

Outbreak Management Team (OMT) - A group of key individuals, including but not limited to, representatives from the Public Health Outbreak Team, Infection Prevention and Control (IPC/ICD), Occupational Health/ Workplace Health Safety (WHS), Facility Administration/Facility Management or their Designate who work cooperatively to ensure a timely and coordinated response to a suspect or confirmed outbreak. Composition of the OMT will depend on disease and facility type.

PPE - Personal Protective Equipment: Equipment worn to minimize exposure to hazards in a workplace.

Provincial Laboratory for Public Health – Also called ProvLab, the provincial lab for Alberta.

Public Health Outbreak Team - For the purpose of this document, the Public Health Outbreak Team coordinates and leads the outbreak response. It also provides consultation and leadership in outbreak investigations in facility settings and is responsible for reporting of facility outbreaks to Alberta Health. This team encompasses the Medical Officer of Health (MOH), and the Communicable Disease Control (CDC) Outbreak Team and Environmental Public Health (EPH) Outbreak Team.

Residents - Tenants who reside in congregate sites.

Respiratory Surveillance Reporting Definition - The threshold to report a potential respiratory outbreak to the Public Health Outbreak Team is called the surveillance reporting definition. See [Table A](#) for further details.

Surveillance Cases - Symptomatic individuals or confirmed cases which are considered indicators of a potential outbreak.

Visitor - An individual who spends time with the resident/patient for a temporary period for the purposes of providing support to the resident/patient and is not an essential partner to care planning and/or decision-making. For more information refer to [AHS Visitor policy](#).

How to Contact the AHS Public Health Outbreak Team to Report Illness

How sites contact the AHS Public Health Outbreak team depends on whether or not initial reporting has already occurred.

Initial point of contact for the Public Health Outbreak Team reporting

AHS Coordinated Early Identification and Response line (CEIR) **1-844-343-0971**

- Acts as the first point of contact for sites in all zones to provide notification to the Public Health Outbreak Team when there are 2 symptomatic cases within 7 days in a congregate setting that **does not** already have a confirmed outbreak.
- If site attempts to contact CEIR outside hours of operation the notification is recommended to be completed the next day

Exception for GI illness: If two residents have **GI symptoms only** and meet the outbreak definition **as outlined in [Section 9](#)** (2 residents meeting GI case definition with onsets within 48 hours of each other), a call is recommended to be made to CEIR at that time to report a potential GI outbreak.

Contacting the Public Health Outbreak Team after site has reported initial cases or an outbreak is declared

Report **additional** information about **respiratory illnesses** (COVID-19, Influenza, and ILI) to the Public Health Outbreak Team (including questions regarding an open respiratory illness outbreak):

- The Facility CDC Outbreak Daily Report Portal (Redcap) is the preferred method to report new positive lab reports and newly symptomatic residents.
- If a site does not have access to the Facility CDC Outbreak Daily Report Portal (Redcap) they are to call the Intake line (instead of calling CEIR) to report newly symptomatic residents and/or new positive lab results.
 - The Intake Line is to be used by all facilities in the North, Edmonton, Central and Calgary Zones for related Respiratory Illness
 - **Intake Line 1-888-522-1919**
 - If facilities in the South Zone do not have access to the Facility CDC Outbreak Daily Report Portal (Redcap) they are to report using Zone specific processes
- If site attempts to report subsequent symptomatic or positive cases outside hours of operation, the notification is recommended to be completed the next day.

Report **additional** information about **Gastrointestinal (GI) Illness** to Environmental Public Health (EPH). The EPH Outbreak Response Lead will provide the site the zone process for reporting subsequent symptomatic residents and/or staff along with zone specific line lists as applicable.

- GI illness **may be** reported using the Facility CDC Outbreak Daily Report Portal (Redcap) or reported by a zone-specific process.

Section 1: Preparing for Potential Outbreaks

It is the responsibility of a facility to always be prepared for the possibility of a respiratory or GI outbreak. Each site is recommended to:

- Ensure staff have access to and are familiar with current AHS outbreak management guides.
- Review and update internal protocols and procedures for outbreak management, including symptoms that require investigation and reporting to the Public Health Outbreak Team ([Table A](#) and [Table B](#)), and case and outbreak definitions for:
 - COVID-19 ([Table E](#))
 - Influenza-like Illness (ILI) ([Table F](#)),
 - Influenza ([Table G](#))
 - Gastrointestinal (GI) illness ([Table H](#))
- Work with key site personnel to ensure adequate availability of supplies for outbreak management including on site specimen kits for respiratory and stool specimen collection.
- The baseline supply inventory that the site requires will depend on the resident/staff number, proximity to the ordering centre and timeframe for supply delivery. Please work with your owner/operator/administration to determine the baseline needed for the site.
- Review routine infection control practices and additional (outbreak management) precautions with staff.
- Ensure staff are aware to contact the AHS CEIR at 1-844-343-0971 when there are 2 symptomatic cases that meet the criteria in either [Table A](#) or [Table B](#) in a congregate setting that do not already have an established outbreak.

Zone Outbreak Preparation Resources

- Refer to local Zone process to access the necessary resources for outbreak preparation
- Long Term Care (LTC) Facility living sites, Designated Supportive Living (DSL), Hospice and Non-designated Supportive Living sites may participate in annual outbreak training through zone specific invitations and processes in the fall.
- [Appendix A](#) contains samples of the documents that facilities may use to prepare for outbreak season.

Roles and Responsibilities for Outbreak Preparation

Note: Due to the complex nature of LTC/DSL/Hospice settings, staffing and resident populations, the individual fulfilling the roles and responsibilities within a site may vary from what is outlined below (e.g., some sites may combine the roles for IPC and WHS, or may have designated staff to fulfill these roles)

The Public Health Outbreak Team (MOH, CDC, EPH)

- Sets the standard of practice for communicable disease surveillance and notification in relation to outbreak investigation and management.
- Develops, maintains, and distributes provincial outbreak resources.
- Provides annual outbreak training to appropriate groups.

Infection Prevention and Control (Infection Control Professional and/or Infection Control Designate)

Note: *In the absence of formal IPC or site ICD coverage, facility administration/manager designates responsibility for these roles. This role may be fulfilled by a single individual at the site, or by a combination of a site Infection Control Designate and supporting groups such as organizational/zone/AHS ICP.*

- Ensures that HCW/staff have access to and are familiar with current AHS outbreak management guide.
- Reviews and updates internal protocols and procedures for outbreak management as necessary, including review of case and outbreak definitions and reporting protocols
- Working with key site personnel, ensures adequate availability of supplies needed for outbreak management including PPE, signage and having unexpired respiratory and stool kits on site for specimen collection as appropriate.
- Reviews routine infection control practices and additional (outbreak management) precautions with HCW/staff

Facility Administration/Facility Management or their Designate

- Supports and encourages the annual seasonal influenza immunization and any recommended COVID-19 immunization of residents, HCW/staff, and volunteers.
 - Refer to [Appendix A](#) - Important Notice to Staff About Influenza and COVID-19
- Develops an influenza response plan that details how antiviral prophylaxis will be implemented for residents (e.g., standing orders) and for HCW/staff (as required).
- Develops a COVID-19 response plan that details how surveillance, isolation, specimen collection, and immunization will be implemented for residents and for HCW/staff.
- Liaises with ICP/ICD to ensure facility/units have protocols and procedures for reporting cases of notifiable diseases and suspected outbreaks as per Section 26 of the [Public Health Act](#)
- Liaises with ICP/ICD or site infection control designate to ensure facility/units have access to a current copy of the AHS outbreak guide and that key HCW/staff (ICP/ICD, front line managers, supervisors, administration on-call, charge nurse, etc.) in the facility have contact information for [reporting outbreaks](#) to the Public Health Outbreak Team via CEIR
- Ensures HCW/staff awareness of symptoms for reporting to CEIR as found in [Table A](#) and [Table B](#)

Site/Unit Manager/Designate

- If required, facilitates DSL residents' Most Responsible Health Practitioner to complete the prescription of advanced treatment for antiviral chemoprophylaxis for use in the event of a confirmed influenza outbreak.
 - In some sites, this is done in collaboration with external partners such as pharmacy.
- In preparation for influenza season, compiles a list of residents (include names, contact and other pertinent information) to be used in the event of an outbreak for implementing antiviral prophylaxis (see [Appendix A](#))
- Liaises with AHS WHS or site OHS designate to ensure HCW have been N95 Respirator fit tested and can use appropriate PPE as dictated by the [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)
- Ensures proper use and adequate supplies of PPE as per [AHS Infection Prevention and Control guidelines for Continuing Care](#)

- Maintains ongoing surveillance and monitoring for unusual clusters of illness in residents and staff (for GI illness) to facilitate early recognition of cases, and identification of a possible outbreak at the site.

Occupational Health/Workplace Health and Safety/Designate

***Note:** In the absence of formal AHS WHS or site OHS coverage, facility administration/manager designates responsibility for these roles.*

- Promotes and provides as appropriate, annual influenza immunization, COVID-19 immunization, and all other recommended immunizations for HCW/staff
- Develops, reviews and updates internal protocols for management of HCW/staff during an outbreak, as necessary.
- Maintains documentation on HCW/staff health and vaccine status and provides the Site/Unit Manager with a list of HCW/staff with reported immunization
- Assesses staff use of PPE and hand hygiene.

Onsite Staff (staff hired directly by the site, contracted staff, AHS staff)

- Ensures they are familiar with current AHS outbreak guide.
- Ensures they are aware of the symptoms for reporting to CEIR as found in [Table A](#) and [Table B](#) as well as COVID-19, ILI, Influenza and GI illness case and outbreak definitions (found in [Section 4](#)) and reporting protocols
- Ensures they are aware of the proper use of PPE as per AHS infection prevention and control guidelines and are aware to use the [Infection Prevention and Control Risk Assessment \(IPCRA\)](#) prior to resident interactions

Section 2: Assessing a Potential Outbreak

2.1 Surveillance and Threshold for Investigation

All sites are responsible for conducting ongoing surveillance and monitoring for unusual clusters of illness in residents and staff (GI only). Surveillance takes place prior to, during and after outbreaks.

Surveillance Cases

Symptomatic individuals or confirmed cases which are considered indicators of a potential outbreak are called “surveillance cases.” The criteria for surveillance cases are found in [Table A](#) and [Table B](#).

- An optional Surveillance Case Tracking Sheet that can be used by sites/facilities can be found in [Attachment 2.1](#).

Respiratory Surveillance Reporting Definition and GI Illness Reporting Definition

The threshold to report a potential outbreak to CEIR is called the respiratory surveillance reporting definition or the GI illness reporting definition.

- The respiratory surveillance reporting definition can be found in [Table A](#).
- The GI illness reporting definition can be found in [Table B](#).

Note: Symptoms not listed in [Table A](#) or [Table B](#) do not need to be reported to the Public Health Outbreak Team. These symptoms may still be indications of illness and are recommended to be managed at the discretion of the care team for the individual experiencing the symptoms.

Table A: Determining when to report respiratory illness symptoms to CEIR

Respiratory Surveillance Case Definition	Respiratory Surveillance Reporting Definition
<p>Residents</p> <p>Residents will count towards the threshold to report symptoms of respiratory illness to CEIR if they develop any of the following:</p> <p>Any new or worsening symptoms listed below:</p> <ul style="list-style-type: none"> • Fever (may not always be present in the elderly) • Cough • Shortness of breath (SOB) • Sore throat • Runny nose/Nasal congestion • Loss of taste and/or loss of smell • Decrease in oxygen (O₂) saturation level or increased O₂ requirements • Nausea or diarrhea 	<p>2 or more surveillance cases in a 7-day period with a common epidemiologic link☆</p> <p>☆<i>Epidemiological link means the cases need to have been in the setting (same site/same unit) during their incubation period or communicable period.</i></p>

Table B: Determining when to report GI illness to CEIR

GI Illness Case Definition	GI Illness Reporting Definition
<p>Residents</p> <p>At least one of the following criteria must be met and not be attributed to another cause (e.g., <i>Clostridioides difficile</i> diarrhea, medication, laxatives, diet, or prior medical condition, etc.):</p> <ul style="list-style-type: none"> 2 or more episodes of diarrhea (i.e., loose, or watery stools) in a 24-hour period, above what is normally expected for that individual <p>OR</p> <ul style="list-style-type: none"> 2 or more episodes of vomiting in a 24-hour period <p>OR</p> <ul style="list-style-type: none"> 1 or more episodes of vomiting AND diarrhea in a 24-hour period <p>OR</p> <ul style="list-style-type: none"> 1 episode of bloody diarrhea <p>OR</p> <ul style="list-style-type: none"> Laboratory confirmation of a known enteric pathogen AND at least one symptom compatible with a GI infection e.g., nausea, vomiting, diarrhea, abdominal pain, or tenderness <p>Notes:</p> <p>Laboratory confirmation is not required.</p> <ul style="list-style-type: none"> If there is an unusual increase in GI illness amongst HCW/staff (above the baseline of what would be expected) whether they were present at work with symptoms or not it is recommended to be reported to CEIR as this could be an indicator of a potential GI outbreak. 	<p>If two residents have GI symptoms only and meet the outbreak definition as outlined in Section 9 (2 residents meeting GI case definition with onsets within 48 hours of each other) a call is recommended to be made to CEIR at that time to report a potential GI outbreak</p>

Stool Specimen Collection

Sites must collect specimens as directed by the Public Health Outbreak Team and arrange for delivery of specimens to the laboratory. It is valuable to collect stool specimens from cases during outbreaks to try and identify the etiology, if possible. Typical GI illness outbreak stool specimen collection procedures are found in [Attachment 3.1](#).

Scenarios which do not meet the Respiratory Surveillance Reporting Definition

- A single resident with symptoms does not meet the respiratory surveillance reporting definition and does not need to be reported to CEIR. Sites must continue to monitor for ongoing illness. An optional Surveillance Case Tracking Sheet can be found in [Attachment 2.1](#).
- Infection control strategies listed in [Figure 2](#) are recommended to be immediately implemented for any resident with symptoms. Prompt implementation of the strategies listed in [Figure 2](#) may help to limit illness from spreading at the site and prevent an outbreak.
- Sites are recommended to update supervisors/managers of one symptomatic resident, so they are aware that a [Surveillance Case Tracking Sheet](#) has been started. Sites are recommended to have an internal policy that dictates how the [Surveillance Case Tracking Sheet](#) (or a site-specific tool) is used to track symptomatic residents and when the call needs to be made to CEIR.

Specimen collection prior to meeting the respiratory surveillance reporting definition

- No specimen collection is required for Public Health Outbreak Team purposes prior to a site meeting the respiratory surveillance reporting definition.
- Sites are recommended to follow internal processes to collect specimens for testing (COVID-19 and RPP) needed for medical management of symptomatic residents as required. Testing may be required to inform interventions such as Oseltamivir (Tamiflu) treatment for influenza cases or [treatment for COVID-19 cases](#) (such as Paxlovid, Remdesivir, Evusheld).
- The Public Health Outbreak Team and CEIR will provide support and guidance on specimen collection recommendations for outbreak management once the respiratory surveillance reporting definition has been met.


Resource - Symptoms? What Do I Do to Prevent a Potential Outbreak?

- See [Appendix B](#)
- The respiratory surveillance reporting definition and the GI illness reporting definition direct sites to call when there are 2 respiratory surveillance cases within 7 days or 2 GI illness cases within 48 hours at the site.
- This resource helps facilities/sites manage symptomatic residents before the reporting definitions are met (that is, prior to meeting the respiratory or GI illness thresholds to call CEIR).
 - This resource may be printed and made available to the front-line staff for daily use.
 - It provides a summary of the steps that are recommended to be taken if the symptomatic person is a resident and who to tell when a resident is symptomatic is also outlined.

2.2 Reporting once the respiratory surveillance or GI illness reporting threshold is met

- Follow the established internal outbreak notification protocols to report residents and HCW/staff (only for GI illness) who display any of the symptoms listed in [Table A](#) and/or [Table B](#) To connect with and report illness to the Public Health Outbreak Team, see section [AHS Zone Public Health Outbreak Team Contacts \(Regular and After Hours\)](#).
- Prompt reporting of a potential outbreak that meets the reporting definitions permits early identification and interventions to interrupt transmission of illness as soon as possible, reducing morbidity and mortality. Initial control measures, directions on recommended specimen collection and PPE recommendations will be provided.
- The Public Health Outbreak Team will follow up with sites after the CEIR notification is received and assess whether sites meet outbreak status. The Public Health Outbreak Team will be responsible for the decision to declare when an outbreak is to be opened.
- Follow internal protocols for site notification about residents that are being tested (e.g., to your IPC/ICD and WHS/OHS designate where available). Also follow the Public Health Outbreak Team instructions for collecting and reporting data if a confirmed outbreak is identified at your site ([Figure 1](#)).
- The information that sites must provide when reporting to CEIR is standardized. See [Table C](#) for a list of all information required.

Figure 1: Potential Outbreak Notification Algorithm*



**This document outlines outbreak reporting in most situations, but there may be variation within some Zones. It is recommended to periodically confirm outbreak reporting procedures within your Zone.*

Table C: Initial Data Required by the Public Health Outbreak Team

Site Information:
Call Received:
Call Returned:
Confirmed Outbreak:
Call Redirected:
Caller Name:
Caller Phone Number:
Caller Email:
Zone:
City:
Site Type:
Is this an AHS Facility:
Site Name:
Site Manager Email:
Site Address:
Postal Code:
Site Phone Number:
Site Fax Number:
Number of Units in Site:
Names of the Unit:
How many residents reside on the unit of the case(s):
Are you able to keep residents to only that area for the duration of the outbreak?
How many staff work on the unit of case(s)?
Are you able to keep staff only to the area for the duration of the outbreak?
Manager Notified at the site?
Total number of residents at the site:
Total number of Staff working at the site:
Additional Notes about the site:
Number of Symptomatic Residents:
Number of Symptomatic Staff:
Do Staff work in multiple locations or on multiple units?
Any Hospitalizations or deaths?

2.3 Reporting Additional or Subsequent Cases to Public Health Outbreak Team

- After the site has made the initial report to CEIR, no further calls to CEIR are required.
 - If the site has not received direction on whether they have a confirmed outbreak, any additional symptomatic residents, and HCW/staff, and/or positive cases (for respiratory illness) are recommended to report using the **Intake Line at 1-888-522-1919** (instead of calling CEIR).
 - If the Public Health Outbreak Team has provided the site with access to report additional symptomatic or positive residents through the Facility CDC Outbreak Daily Report Portal (Redcap), additional symptomatic residents may be reported here instead of reporting through the Intake Line.
- For South Zone, the intake line is NOT used to report additional cases of illness. Zone specific processes are used.

- Report additional **Gastrointestinal** Illness cases that meet GI case definition during a GI outbreak to Environmental Public Health. The EPH Outbreak Response Lead will provide the site the zone process for reporting subsequent symptomatic residents and/or staff along with zone specific line lists as applicable.

Roles and Responsibilities: Assessing for and Reporting Potential Outbreaks

Note: Due to the complex nature of LTC/DSL/Hospice settings, staffing and resident populations, the individual fulfilling the roles and responsibilities within a site may vary from what is outlined below (e.g., some sites may combine the roles of IPC and WHS, or may have designated staff to fulfill these roles)

The Public Health Outbreak Team (MOH, CDC, EPH)

- Provides consultation on suspected clusters of illness or outbreaks.
- Determines the need to initiate an outbreak investigation.
- Requests information from their designated contact at the Facility (e.g., site Administrator, IPC, ICD) to meet the Public Health Outbreak Team requirements for outbreak management.

Infection Prevention and Control (Infection Control Professional and/or Infection Control Designate)

Note: In the absence of formal IPC or site ICD coverage, facility administration/manager designates responsibility for these roles. This role may be fulfilled by a single individual at the site, or by a combination of a site Infection Control Designate and supporting groups such as organizational/zone/AHS ICP.

- Acts as a resource for frontline HCW/staff to facilitate early recognition of possible outbreaks.
- Notifies the AHS CEIR line as soon as possible in the event of a suspected outbreak as per site process - see [Figure 1](#).
 - In facilities where there is no one assigned the role of IPC, the front line HCW/staff or manager at the outbreak unit/site notifies the AHS CEIR line.
- Obtains reports on the clinical status of all affected individuals and works with HCW/staff to identify new cases.

Facility Administration/Facility Management or their Designate

- Responsible for maintaining operations to provide optimal care and services for symptomatic residents.
- Notifies senior management within the site as indicated by internal protocols.
- Notifies IPC/ICD when an unusual cluster of illness is suspected. In facilities where there is no one assigned the role of IPC, contact the AHS CEIR line - see [Figure 1](#)
- Directs the immediate implementation of routine IPC measures.
- Communicates resident assessment, monitoring, surveillance, and reporting to the Public Health Outbreak Team and IPC/ICD as appropriate.

Onsite Staff (staff hired directly by the site, contracted staff, AHS staff)


- Place symptomatic residents on appropriate precautions and ensure all signs and symptoms of resident illness are documented as per site standards.
- Assesses that illness in symptomatic residents meets the reporting criteria in [Table A or Table B](#)
- Notifies the AHS CEIR line promptly if assigned when the reporting definition has been met.

- Works collaboratively with all partners to ensure ongoing monitoring and surveillance at the site to identify new case defining illness in residents.
- Liaises with Most Responsible Health Practitioner and external partners, as necessary.
- Provides services to care and treat symptomatic residents in place as per [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)

Attachment 2.1: Surveillance Case Tracking Sheet

This tool can help sites keep track of symptomatic residents to determine if the site is recommended to call CEIR. Sites are recommended to update supervisors/managers of one symptomatic resident or HCW/staff such that they are aware that a Surveillance Case Tracking Sheet has been started. Sites are recommended to have an internal policy that dictates how the Surveillance Case Tracking Sheet (or a site-specific tool) is used to track symptomatic residents or HCW/staff and when the call needs to be made to CEIR.

This resource can be printed and made available to the front-line staff for daily use. Please print copies directly from the web page online - [Outbreak Management | Alberta Health Services](#)



**Alberta Health
Services**

Surveillance Case Tracking Sheet

Name (Last Name, First name)	Resident/ Staff	Onset of Symptoms (dd-Mon-yyyy)	Symptoms	For Staff Cases - Date Worked While Symptomatic
	<input type="checkbox"/> Resident <input type="checkbox"/> Staff		<input type="checkbox"/> Fever <input type="checkbox"/> Nausea/vomiting/diarrhea Any new or worsening symptoms listed below: <input type="checkbox"/> Cough <input type="checkbox"/> Shortness of breath (SOB) <input type="checkbox"/> Sore throat <input type="checkbox"/> Runny nose/nasal congestion <input type="checkbox"/> Loss of taste and/or loss of smell <input type="checkbox"/> Decrease in oxygen (O ₂) saturation level or increased O ₂ requirement	
	<input type="checkbox"/> Resident <input type="checkbox"/> Staff		<input type="checkbox"/> Fever <input type="checkbox"/> Nausea/vomiting/diarrhea Any new or worsening symptoms listed below: <input type="checkbox"/> Cough <input type="checkbox"/> Shortness of breath (SOB) <input type="checkbox"/> Sore throat <input type="checkbox"/> Runny nose/nasal congestion <input type="checkbox"/> Loss of taste and/or loss of smell <input type="checkbox"/> Decrease in oxygen (O ₂) saturation level or increased O ₂ requirement	
	<input type="checkbox"/> Resident <input type="checkbox"/> Staff		<input type="checkbox"/> Fever <input type="checkbox"/> Nausea/vomiting/diarrhea Any new or worsening symptoms listed below: <input type="checkbox"/> Cough <input type="checkbox"/> Shortness of breath (SOB) <input type="checkbox"/> Sore throat <input type="checkbox"/> Runny nose/nasal congestion <input type="checkbox"/> Loss of taste and/or loss of smell <input type="checkbox"/> Decrease in oxygen (O ₂) saturation level or increased O ₂ requirement	
	<input type="checkbox"/> Resident <input type="checkbox"/> Staff		<input type="checkbox"/> Fever <input type="checkbox"/> Nausea/vomiting/diarrhea Any new or worsening symptoms listed below: <input type="checkbox"/> Cough <input type="checkbox"/> Shortness of breath (SOB) <input type="checkbox"/> Sore throat <input type="checkbox"/> Runny nose/nasal congestion <input type="checkbox"/> Loss of taste and/or loss of smell <input type="checkbox"/> Decrease in oxygen (O ₂) saturation level or increased O ₂ requirement	
	<input type="checkbox"/> Resident <input type="checkbox"/> Staff		<input type="checkbox"/> Fever <input type="checkbox"/> Nausea/vomiting/diarrhea Any new or worsening symptoms listed below: <input type="checkbox"/> Cough <input type="checkbox"/> Shortness of breath (SOB) <input type="checkbox"/> Sore throat <input type="checkbox"/> Runny nose/nasal congestion <input type="checkbox"/> Loss of taste and/or loss of smell <input type="checkbox"/> Decrease in oxygen (O ₂) saturation level or increased O ₂ requirement	

This resource is intended to be used as part of the surveillance and assessment strategies outlined in the provincial outbreak Guides. Details can be found in the **Guide for Outbreak Prevention & Control in Non-Designated Supportive Living (NDSL) Sites** and the **Guide for Outbreak Prevention & Control in Long Term Care and Designated Supportive Living Sites**

22017(2022-06v2)

Section 3: Managing a Potential Outbreak

Sites that have 2 or more cases and meet the Reporting Definition listed in [Table A](#) or [Table B](#) will receive further direction from CEIR.

- This direction will include information on IPC measures, specimen collection, and outbreak control strategies.
- The Public Health Outbreak Team will review the report from the site and will advise the site if they meet either the respiratory or GI outbreak reporting definition and have a potential or confirmed outbreak.
- Staff, IPC/ICD, and the Public Health Outbreak Team in congregate settings work collaboratively with facility administrators and HCW/staff to facilitate prompt response to help minimize the impact of the outbreak. For ongoing updates relevant to congregate settings, log in to [Continuing Care Connection](#).

Note: You will be required to register the first time you use the site.

3.1 Infection Prevention and Control Measures

- In the event a single resident develops symptoms in accordance with [Table A](#) or [Table B](#) the IPC Practices and Additional Precautions in [Figure 2](#) shall be implemented immediately. Do not wait for the causative agent to be identified.
- Isolation of residents with symptoms:
 - Symptomatic residents
 - Isolate immediately using additional precautions. Refer to [Figure 2](#). Staff to wear appropriate PPE as determined by [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)
 - Place signage outside the resident's room, near the door, alerting HCW/staff and visitors that the resident is symptomatic, and precautions are required.
- Strict hand hygiene is the most important measure in preventing spread of infections for both HCW/staff and residents.
 - Hand hygiene is recommended to be performed in accordance with the [AHS Hand Hygiene Policy and Procedure](#) that provides direction on product selection, location, and use.
 - Alcohol-based hand rubs containing a minimum of 60-90% alcohol are as effective as soap and water when hands are not visibly dirty. They are recommended to be clearly labelled with a DIN or label claim as being effective and used prior to expiry date.
 - Wash hands with soap and water when:
 - Hands are visibly soiled with food, dirt or blood and body fluids.
 - Prior to and during and after handling food
 - Immediately after using washroom facilities.
 - Following glove removal (doffing)
 - Glove use is not a substitute for hand hygiene; hand hygiene is required after glove removal.
- **Routine practices** help prevent the spread of infection and reduce the possibility that HCW/staff will sustain accidental exposures to infectious organisms. Routine practices are used for every resident, every time regardless of their diagnosis or infectious status.
- **Additional precautions** such as droplet and contact precautions are determined and implemented based on presenting symptoms using the [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)

Figure 2: Routine Practices and Additional Precautions

Precautions for Respiratory and GI Illness

All HCW/staff must conduct an [Infection Prevention and Control Risk Assessment \(IPCRA\)](#) prior to every resident interaction – check [Infection Prevention & Control | Alberta Health Services](#) for the most current recommendations.

Implement recommended precautions ([Modified Respiratory Precautions](#) and [Droplet and Contact Precautions](#)) when caring for symptomatic residents to control the spread of viruses.

Precautions Posters available at [Posters | Alberta Health Services](#) and [Routine practices](#) in Continuing Care for further information.

Resident Placement and Signage

- Single room preferred.
- Maintain a distance of 2 metres between residents sharing a room use of physical barriers (curtains or portable wipeable screens) is recommended to be implemented as all times.

Mask

- Wear appropriate mask for any encounter, within two (2) metres, with a resident who has, or is suspected of having a respiratory illness. Use the mask (surgical, KN95, N95) recommended by the PCRA.
 - N95 Respirator (fit-tested) - for aerosol generating medical procedures (AGMP) and residents on airborne precautions.
 - AGMPs are defined as any medical procedure that can induce the production of aerosols of various sizes, including droplet nuclei. [See Aerosol Generating Medical Procedures.](#)

Eye Protection

- When a mask or N95 respirator is worn, eye protection or face shields are recommended to be worn as indicated by the PCRA or additional precautions.
- Personal (prescription) eyewear does not provide adequate protection.

Gowns

- Gowns are used to protect clothing and forearms from contamination of droplets and from direct contact with residents or residents' environment.

Gloves

- Wear clean non-sterile gloves for direct contact with resident or resident's environment and to protect from droplet exposure.

Hand Hygiene (4 moments from [AHS Hand Hygiene Policy and Procedure](#))

- Before contact with a resident or resident's environment including but not limited to putting on (donning) personal protective equipment; before entering a resident's room; and, before providing resident care.
- Before a clean or aseptic procedure including but not limited to wound care; handling intravenous devices; handling food; or preparing medications.
- After exposure (or risk of exposure) to blood and/or body fluids including but not limited to when hands are visibly dirty, following removal of gloves.
- After contact with a resident or resident's environment including but not limited to removing (doffing) personal protective equipment, leaving a resident's environments and after handling resident care equipment.

Resident Care Equipment

- Dedicate to single resident or clean and disinfect after each use.

Resident Transport

- Transport for essential purposes only.
 - Residents wear an appropriate mask during transport (if tolerated).
 - Notify receiving department that resident is on additional precautions.
- *If resident is confirmed to have a GI pathogen refer to [Section 9](#) for further information

AHS PPE Donning and Doffing posters available at: [COVID-19 Personal Protective Equipment \(PPE\) | Alberta Health Services](#)

Visitors: Request visitors report to administration desk or nursing desk to discuss precautions before entering resident's room.

Environmental Services: change mop head, cloths, and cleaning solution after cleaning each room or bed space.

The [Infection Prevention and Control in Continuing Care](#) webpage has numerous additional resources.

3.2 Administrative Measures

- Ensure adequate availability of all supplies (e.g., hand hygiene products, PPE, linen, specimen collection supplies) through notification of appropriate departments.
- Ensure HCW/staff are maintaining heightened surveillance to identify and report newly symptomatic residents as per [Table A](#) and [Table B](#).
- Consult with the ICP/ICD or the Public Health Outbreak Team when making decisions about cohorting HCW/staff assignments. Cohorting is recommended where operationally feasible (i.e., symptomatic residents receiving treatments after asymptomatic residents, or having designated HCW/staff treat symptomatic clients). Consider:
 - Cohorting HCW/staff to affected areas if practical or assigning HCW/staff to care for asymptomatic residents before symptomatic residents.
 - Minimizing movement of HCW/staff, students, or volunteers between floors/areas, especially if some areas are not affected.
 - Cohorting residents with the same illness
 - Cohorting exposed asymptomatic residents.

3.3 Resident Restrictions

Note: *If necessary, consult with IPC/ICD for assistance with adapting resident activities.*

- Asymptomatic residents and those who are not a case are able to participate in daily activities.
- Symptomatic residents are recommended to remain in their rooms. Additional precautions are required when entering a symptomatic resident's room (See [Figure 2](#)).
 - If a symptomatic resident must leave their room, they are recommended to wear an appropriate mask (as tolerated).
- Symptomatic residents are not recommended to participate in group activities.
- During a potential outbreak, symptomatic residents are recommended to only receive treatment such as physiotherapy or occupational therapy in their rooms instead of a centralized area. They are recommended to wear an appropriate mask (as tolerated).
- Symptomatic residents will be allowed to attend medically necessary activities or appointments, and they are recommended to wear an appropriate mask (as tolerated). Ensure receiving facility/unit is notified of the potential outbreak so that appropriate precautions can be taken for the resident on arrival.
- If resident chooses not to wear a mask, HCW is recommended to review their PCRA and adjust PPE accordingly (e.g., apply modified respiratory precautions until a thorough symptom and risk factor history can be obtained).
- For symptomatic residents, if COVID-19 or influenza is suspected, see early treatment recommendations in [Section 6.2](#) and [Section 8.2](#) respectively.
- Additional considerations for accommodations that may be supported for safe movement of residents with dementia or cognitive impairment who are in isolation:
 - Support the resident to leave their room only in ways that minimize spread of infection (e.g., one-on-one support to the resident at all times when they are out of their room, putting on PPE, using hand sanitizer, avoiding others, and avoid touching surfaces, etc.); and
 - Minimize contact with the isolated resident (e.g., minimize the possibility of other residents going into that person's room) by offering additional activities and interventions for non-isolated residents in the unit.
 - Support outdoor visits with DFSP and volunteers.
 - Encourage activities that engage and support the resident without resulting in risk to others.

3.4 Restrictions on Affected Unit/Site

- No restrictions would be implemented at this point in the investigation.

3.5 Admissions/Transfers from Acute Care Site to an LTC, DSL or Hospice Site

- No changes from usual practices would be implemented at this point in the investigation.

3.6 Transfers from an Outbreak LTC, DSL or Hospice Site to an Acute Care Site

- If a symptomatic resident requires transfer to an acute care site, the facility must notify the EMS* Dispatcher, the transport staff (EMS crew) and the receiving care facility that the resident is under investigation and appropriate PPE must be maintained.

* In some communities, [EMS Community Paramedics](#) can provide immediate or scheduled medical supervision and treatments that are currently unavailable in the community setting. A referral is required to access this service.

3.7 Group/Social Activities and Other Events

- Symptomatic residents are not recommended to participate in group or social activities with other residents.
 - Symptomatic residents may continue to participate in social activities with visitors and DFSPs. This includes going outdoors and participating in 1:1 activities as long as additional precautions are followed such as enhanced hand hygiene, masking and physical distancing.
- No other changes to group or social activities are required at this point in the investigation.

3.8 Nourishment Areas/Sharing of Food

- Symptomatic residents are recommended to receive meal service in their room.

3.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3)

- No changes from usual practice would be recommended by the Public Health Outbreak Team at this time.

3.10 Visitors and Designated Family/Support Person(s) (DFSPs)

- Request that all visitors/DFSPs report to the reception/nursing desk before visiting residents.
- Visitors/DFSPs are recommended to be advised of potential risk of exposure if visiting a symptomatic resident.
- Symptomatic visitors/DFSPs are not recommended to visit.
- Recommend visitors/DFSPs wear a mask while at the facility due to potential outbreak.
- In extenuating circumstances, site will determine if visitation is recommended when a visitor/DFSP is symptomatic.
 - Compassionate exemptions support visitation by symptomatic visitors/DFSPs of patients who are at end of life.
- Advise visitors/DFSPs visiting symptomatic residents to wear PPE (e.g., gloves, gown, appropriate mask, eye protection) and to clean hands with alcohol-based hand rub before putting on and removing the PPE.
- Demonstrate for visitors/DFSPs how to utilize PPE appropriately.

- Request visitors/DFSPs to follow the directions of HCW/staff and Facility Administration.

Designated Family/Support Person(s)

- DFSPs are essential partners in the provision of safe, quality resident care.
- Facilities are recommended to support the presence of residents DFSPs while balancing the safety of all residents, DFSPs, visitors and HCW/staff.
- Follow [Visiting a Loved One at an AHS Facility](#) (or for non-AHS sites, follow site policy regarding DFSPs)
- DFSPs are not recommended to be restricted from visits, but limits may be required. Refer to [Visiting a Loved One at an AHS Facility](#) (or for non-AHS sites, follow site policy regarding limits for DFSP visits).

3.11 Volunteers

- Advise volunteers of the importance of hand hygiene and routine precaution protocols.
- Symptomatic volunteers are not recommended to visit the site.
- Volunteers are recommended to follow the same PPE recommendations as HCW/staff (see [3.12](#)).

3.12 HCW/Staff Outbreak Control Measures (Including volunteers, students, and physicians)

- [Table D](#) outlines HCW at home rapid antigen testing recommendations and work restrictions provided below.
- HCW/staff are recommended to monitor themselves for signs and symptoms of illness.
- Symptomatic HCW/staff are recommended to take an at-home rapid antigen test. [See Table D](#). Symptomatic HCW/staff are not recommended to report to work. They are recommended to report illness to their manager/designate.
 - They are recommended to report to WHS/OHS if symptoms are related to workplace exposure.
- HCW/staff who develops respiratory symptoms at work are recommended to perform respiratory hygiene practices (e.g., coughing into sleeve, using tissues, wearing an appropriate mask) and leave the workplace as soon as possible. HCW/staff who develops GI symptoms at work are recommended to perform hand hygiene and leave the workplace as soon as possible.
- Cohort or assign HCW/staff to care for asymptomatic residents before symptomatic residents when possible.
- Consider minimizing movement of HCW/staff, students, or volunteers between units/floors, especially if some units are not affected.
If possible, during initial investigation, assign HCW/staff that have been immunized against influenza and COVID-19 to care for symptomatic residents.
- Ideally, HCW/staff assigned to housekeeping duties are not recommended to be involved in food preparation or food service during outbreaks. If this is not possible, ensure meticulous attention is paid to IPC practices, and consider completing any required food preparation before beginning housekeeping.

Note: AHS and Covenant Health HCW/staff are required to follow all their organizations policies and procedures regarding work attendance, masking and eye protection.

Table D: Health Care Worker/Staff Testing and Work Restrictions

HCW/staff Rapid Antigen Testing Recommendations and Resulting Work Restrictions	
<p>Symptomatic refers to new onset of fever, cough, shortness of breath, sore throat, runny nose/nasal congestion, loss of taste and/or loss of smell, nausea/diarrhea.</p> <p>HCW/staff who are experiencing the above symptoms are recommended to complete a Rapid Antigen Test</p>	
Scenarios	Recommendations
Scenario A Symptomatic with initial negative Rapid Antigen Test	<ul style="list-style-type: none"> HCW/staff is recommended to complete a second Rapid Antigen Test at least 24 hours from the first Rapid Antigen Test If completing a second Rapid Antigen Test: <ul style="list-style-type: none"> Second Rapid Antigen Test POSITIVE: Follow directions in Scenario B below. Second Rapid Antigen Test NEGATIVE: continue to stay home until symptoms improve/are fever-free for 24 hours without use of fever reducing medication, whichever is longer and well enough to resume normal activities. When the HCW returns to work, perform frequent and careful hand hygiene. A mask is recommended to be worn at all times for a total of 10 days from the onset of symptoms. Masking is not necessary if the HCW/staff already isolated for 10 days or more. If not completing a second Rapid Antigen Test, follow directions in Scenario C below.
Scenario B Symptomatic with positive Rapid Antigen Test	<ul style="list-style-type: none"> Work restricted for at least 5 days or until symptoms improve and fever free for 24 hours without use of fever reducing medication (whichever is longer). When the HCW returns to work a mask is recommended to be worn at all times for a total of 10 days from the onset of symptoms (even if the symptoms have improved or resolved). Perform frequent and careful hand hygiene. Masking is not necessary if HCW/staff already isolated for 10 days or more.
Scenario C Symptomatic HCW/staff (no Rapid Antigen Test or ONLY one Negative Rapid Antigen Test)	<ul style="list-style-type: none"> Encourage HCW/staff to complete Rapid Antigen testing. If testing is not completed: Work restricted for at least 5 days or until symptoms improve and fever free for 24 hours without use of fever reducing medication (whichever is longer). Once work restriction is complete, When the HCW returns to work a mask is recommended to be worn at all times for a total of 10 days from the onset of symptoms (even if the symptoms have improved or resolved). Perform frequent and careful hand hygiene. Masking is not necessary if HCW/staff already isolated for 10 days or more.
Scenario D Symptomatic HCW/staff Symptoms NOT listed above.	<ul style="list-style-type: none"> Rapid Antigen Testing not recommended. Continue to stay home until symptoms improve and well enough to resume normal activities.

3.13 Specimen Collection

Not all residents will be recommended to have specimens collected for outbreak management.

- Once the respiratory surveillance or GI illness reporting definition has been met (see [Table A](#) and [Table B](#)) collect the appropriate specimen samples:
 - **Respiratory illness:** symptomatic residents is recommended to have a NP swab collected to determine the causative agent.
 - The number of residents who need specimens collected will be determined in consultation with the Public Health Outbreak Team. The Public Health Outbreak Team will advise the site of exactly what testing (e.g., COVID-19, Influenza, RPP) is recommended to be ordered.
 - **GI illness:** collect stool specimen samples only on the direction of the Public Health Outbreak Team or MOH.
- If specimen collection is required by the Public Health Outbreak Team, ensure proper collection and labelling of specimens (including using assigned EI number on all specimens).
- Make internal arrangements for transporting specimens to the lab. See ProvLab Specimen Collection and Transport Guidance ([Attachment 3.1](#)).
- Specimen collection is recommended to be ordered by the Most Responsible Health Practitioner and not under the Zone MOH.

3.14 Enhanced Environmental Cleaning and Disinfection

- Clean and disinfect:
 - Common/public areas
 - At least once per day for low touch surfaces (e.g., shelves, benches, windowsills, message, or white boards, etc.),
 - a minimum of two times daily for high touch surfaces (e.g., doorknobs, light switches, call bells, handrails, phones, elevator buttons, TV remote), care/treatment area, dining areas and lounges.
 - Immediately on any visibly dirty surfaces.
 - Use a “wipe twice” procedure (a 2-step process) to clean and then disinfect surfaces (i.e., wipe surfaces thoroughly to clean visible dirt then wipe again with a clean cloth saturated with disinfectant to disinfect) while observing the appropriate contact time.
 - Health care equipment (e.g., wheelchairs, walkers, lifts) is recommended to be cleaned and disinfected according to manufacturer’s instructions.
 - Any shared resident health care equipment (e.g., commodes, blood pressure cuffs, thermometers, lifts, bathtubs, showers, shared bathrooms) is recommended to be cleaned and disinfected after use and prior to use by a different resident.
 - All HCW/staff equipment (e.g., computer keyboards/mouse/carts and/or screens, medication carts, charting desks or tables, telephones, touch screens, chair arms) is recommended to be cleaned and disinfected at least twice daily and when visibly dirty.
 - Refer to manufacturer instructions for cleaning and disinfection procedures, including compatible cleaners and disinfectants.
- Areas that are not considered common/public areas (e.g., private offices administrative areas, etc.) do not require enhanced cleaning/disinfection.
- Operators of facilities may develop an approach to environmental cleaning and disinfection that includes their HCW/staff, service providers (e.g., home care), designated family/support persons and visitors.

- If the surface disinfectant product used has cleaning properties (detergent/disinfectant), it may be used for both steps. Follow manufacturer's directions for use.
- Use a disinfectant with a Drug Identification Number (DIN) and broad spectrum virucidal claim, or a specific virucidal claim against non-enveloped viruses and coronaviruses for enhanced general environmental cleaning.
- Be sure to use the appropriate precautions when using chemicals for cleaning and disinfecting. Consult the products Material Safety Data Sheets. Perform cleaning using the proper personal protective equipment (PPE). Follow correct donning and doffing of PPE.
 - [Putting on \(Donning\) Personal Protective Equipment](#) and [Taking off \(Doffing\) Personal Protective Equipment \(PPE\)](#) instructions on the AHS IPC.
- Upholstered furniture, rugs or carpets contaminated with bodily fluid (emesis or stool) are difficult to clean and disinfect properly. Consult manufacturer's instructions for the cleaning instructions. If not available, remove the item from the resident area and steam clean as soon as possible. Consider discarding items that cannot be appropriately cleaned/disinfected, when possible/appropriate.
- HCW/staff handling soiled laundry is recommended to wear gloves, and gowns if there is a risk of contaminating clothing.
- If the laundry machine has been used to clean laundry soiled with diarrhea or vomiting, a bleach cycle of the laundry machine is recommended to be run (without a load of laundry) before washing other laundry.

Specific information about environmental cleaning can be found at the links below:

- [Principles for Environmental Cleaning and Disinfection \(albertahealthservices.ca\)](#)
- [Public Health Recommendations for Environmental Cleaning and Disinfection of Public Facilities](#)

Roles and Responsibilities for Managing a Potential Outbreak

Note: Due to the complex nature of LTC/DSL/Hospice settings, staffing and resident populations, the individual fulfilling the roles and responsibilities within a site may vary from what is outlined below (e.g., some sites may combine the roles or IPC and WHS, or may have designated staff to fulfill these roles)

The Public Health Outbreak Team (MOH, CDC, EPH)

- Facilitates laboratory testing by recommending type of specimen to be collected and testing required.
- Recommends best practice outbreak control measures to be implemented including admission/transfer status (open or restricted), immunization, chemoprophylaxis if appropriate, and management of HCW/staff.
- Reviews daily illness reporting received from site, monitors potential outbreak progress, and provides consultation to the site when appropriate.
- Obtains or opens an Exposure Investigation number (EI number) through the ProvLab (if not already opened by CEIR) for the tracking of all outbreak-related specimens and communicates the EI number to the site.

Infection Prevention and Control (Infection Control Professional and/or Infection Control Designate)

Note: *In the absence of formal IPC or site ICD coverage, facility administration/manager designates responsibility for these roles. This role may be fulfilled by a single individual at the site, or by a combination of a site Infection Control Designate and supporting groups such as organizational/zone/AHS ICP.*

- Provides IPC recommendations for site to implement to prevent further transmission.
- Directs the implementation of initial IPC measures immediately; does not wait until the etiology is confirmed.
- Obtains reports on the clinical status of all affected individuals and works with HCW/staff to identify new cases.
- Provides the Public Health Outbreak Team with status updates of relevant changes to the investigation such as additional cases, hospitalizations, or deaths.
 - The Public Health Outbreak Team will provide the method on how this information is recommended to be reported.
- Provides resident details to OHS/WHS for possible staff close contact identifications as required for outbreak management.
- In collaboration with the Public Health Outbreak Team, coordinates the collection of clinical specimens as appropriate and provides instruction on completion of requisitions, including the EI number.
- Completes site reviews/audits as needed.

Facility Administration/Facility Management or their Designate

- Responsible for maintaining operations to provide optimal care and services for symptomatic residents.
- Collaborates with IPC/ICD and the Public Health Outbreak Team on control strategies.
- Works collaboratively with the Public Health Outbreak Team to disseminate information to HCW/staff, residents, students, other professions working in the facility that are not HCW, other departments and families, as needed.
- Communicates with HCW/staff, residents, families, volunteers, students, and visitors within the facility as appropriate.
- Consult OHS/WHS and/or IPC/ICD as needed.
- Communicates to appropriate stakeholders outside of the facility.
- Communicates resident assessment, monitoring, surveillance, and reporting to the Public Health Outbreak Team and/or IPC/ICD.
- Collaborates with WHS/OHS designate to identify and report symptomatic HCW/staff to the Public Health Outbreak Team if appropriate.
- Will act as a contact for HCW/staff calling in ill and advise not to attend work.
- Anticipates and provides adequate unit resources for potential outbreak management.
- Ensures proper collection of appropriate specimens as directed by the Public Health Outbreak Team, including using assigned EI number on all specimens, and makes own internal arrangements for transporting specimens to the lab.
 - Testing is recommended to be ordered by the Most Responsible Health Practitioner and not under the Zone MOH
- Maintains clear and consistent channels of communication within the potential outbreak site as per established zone practices.
- Ensures proper use and adequate supplies of PPE as per AHS infection prevention and control guidelines.

Occupational Health/Workplace Health and Safety/Designate

Note: *In the absence of formal AHS WHS or site OHS coverage, facility administration/manager designates responsibility for these roles.*

- Maintains close communication with Frontline Unit/Site Manager with respect to HCW/staff work restrictions/return to work assessments as applicable.
- Provides information to HCW/staff about work restrictions.
 - See [Section 6.12](#) for COVID-19, [Section 8.12](#) for Influenza-specific and [Section 9.12](#) for GI recommendations
- Follows recommendations as directed by the Public Health Outbreak Team and/or WHS/OHS department.
- Supports illness assessment and surveillance of HCW/staff from potential outbreak unit.
- Reports HCW/staff GI illness directly to the Public Health Outbreak Team or provides the information to site designate for the Public Health Outbreak Team reporting.
- Assesses for HCW/staff PPE breaches or hand hygiene breaches as appropriate.

Onsite HCW/Staff (staff hired directly by the site, contracted staff, AHS staff)

- Works collaboratively to facilitate outbreak investigations and implement appropriate initial infection control measures immediately.
 - It is not necessary to wait until the etiology is confirmed.
- Coordinates the collection of clinical specimens as appropriate, under the direction of the Public Health Outbreak Team.
- Works collaboratively to ensure ongoing monitoring and surveillance at the site to identify newly symptomatic illness in residents according to [Table A](#) or [Table B](#).
- Maintains clear and consistent channels of communication within the potential outbreak site as per established Zone practices.
- Liaises with Most Responsible Health Practitioner, as necessary.
- Provides services to care and treat residents in place as per [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)

Provincial Laboratory for Public Health (ProvLab)

- Designates laboratory contact (e.g., microbiologist or virologist) for each outbreak, as required.
- Assigns EI number to facilitate specimen tracking in cases where the Public Health Outbreak Team has not opened the EI independently (applies only to respiratory surveillance purposes only)
- As required, provides consultation to the Public Health Outbreak Team on specimen type and testing appropriate for the outbreak, including genotyping
- Provides specimen collection supplies, as required.
- Ensures the Public Health Outbreak Team and IPC/ICD (if noted on the requisition) receive timely results of potential outbreak specimens.
- Tracks all samples submitted under the EI number.

Attachment 3.1: ProvLab Specimen Collection Guidance

Check ProvLab Bulletins for most current information on specimen collection, testing, and interpretation of lab results.

[Public Health Laboratory \(ProvLab\)](#) or [Forms & Requisitions | Alberta Health Services](#)

Instructions and demonstrations for collection of various types of specimens, including nasopharyngeal swabs can be accessed through the AHS ProvLab website:

[Education Resources | Alberta Health Services](#)

The Laboratory Policy for Acceptance of Laboratory Samples, Test Directories, TDG and other collection information can be found on the AHS ProvLab website:

[Laboratory Test Directory & Collection Information | Alberta Health Services](#)

The specimen requisition must be completed to include:

- Resident's full name (first and last names)
- Resident Personal Health Number (PHN) or unique numerical assigned equivalent
- Resident demographics including date of birth (DOB), gender, address, phone number
- Most Responsible Health Practitioner name (full name), address/location
- Test orders clearly indicated, including body site and sample type, date, and time of collection
- Clinical history and other clinical information
- Facility name, and if applicable, unit
- EI number (assigned by CEIR/ ProvLab/the Public Health Outbreak Team)
- Fax number of outbreak facility/unit or ICP/ICD office
 - Results will be faxed to the outbreak facility/unit or ICP/ICD **when it is noted on the requisition.**

Nasopharyngeal (Np) and Throat Swab for Detection of Respiratory Infections

General Information:

- Unless otherwise directed by the Public Health Outbreak Team, symptomatic residents are recommended to be tested for both COVID-19 and RPP.
- The amount of virus is greatest in acute phase of illness, usually within the first 48-72 hours of symptom onset.
- NP swabs are the preferred specimens for respiratory virus testing. See [ProvLab education resources](#) for information on collection of NP and Throat swabs.
 - If nasopharyngeal swabs are difficult to collect, throat swabs collected in viral transport media are acceptable alternatives for COVID testing. An RPP cannot be completed on a throat swab.
- Once an etiologic agent has been identified, follow the Public Health Outbreak Team direction on the type of testing required for subsequent symptomatic residents and HCW/staff as appropriate.

Stool Specimen Information

- Sites must collect specimens as directed by the Public Health Outbreak Team and arrange for delivery of specimens to the laboratory. It is valuable to collect stool specimens from cases during outbreaks to try and identify the etiology, if possible.

- Please note that norovirus can be detected by ordering Gastro Viral Panel (GVP) if in stool but cannot presently be isolated from vomitus, therefore the collection of vomitus specimens is not recommended for GI illness outbreak management.
- A unique EI number is assigned to each specific outbreak. The Public Health Outbreak Team will obtain an EI number from the ProvLab when a GI illness outbreak is declared. Stool specimens submitted without an EI number on the requisition may not be analyzed for norovirus (i.e.); therefore, it is important that an EI number be obtained prior to collection of outbreak stool specimens.
- The typical turnaround time for norovirus PCR results from the ProvLab (i.e., time between receipt of the specimen at the lab and report of results) is 48 hours. Results are also available on Netcare within 48 hours. The Public Health Outbreak Team will report the result to the ICP/ICD/designate within one business day of receipt of results from the lab.

Procedures to collect stool specimens

- As directed by the Public Health Outbreak Team, collect stool specimens from residents that are acutely ill with diarrhea, preferably within 24-48 hours of onset of symptoms.
- Collect one stool specimen from up to 5 symptomatic residents per outbreak investigation (EI number), preferably during the acute phase of illness. This number of specimens is usually sufficient to determine the etiology of the outbreak.
- Collect stool in a specimen collection “hat” or other clean and dry receptacle (e.g., bed pan, margarine container).
- Do not mix stool with urine or water.
- Place the stool in a clean dry specimen container by using a scoop from stool collection kit, or a disposable tongue depressor or plastic spoon, keeping the outside of the container clean. Fill the container with stool up to one third or at least one tablespoon full and discard the remaining stool. (Sterile container may include container from stool collection kit or sterile urine container).
- Screw the lid tightly to avoid leakage.
- Put the container with the stool into the plastic (biohazard) bag and seal the bag.
- Complete the ProvLab requisition form to include the EI# and the resident’s full first and last names; Personal Health Number (PHN) or unique numerical assigned equivalent; resident demographics to include date of birth (DOB), gender, address, phone number; physician full name and complete address/location; test orders clearly specified including body site and sample type; date and time of collection.
- Label the sample container with the EI number, resident’s full first and last names, PHN or unique numerical equivalent, and date of sample collection.
- Keep stool specimens in the fridge (not the freezer) until ready for transport.
- Batch specimens together and transport to the ProvLab within 24 hours.
- If one or more of these samples are positive and an etiological agent has been identified, then further specimens are not recommended to be collected unless advised by the Public Health Outbreak Team. If additional specimens are received under the same EI# at some later period, these will not be tested unless the Public Health Outbreak Team has contacted the ProvLab point person for the EI number (e.g., MOC/VOC/Designate).
- If all batched samples received have been tested and if all are negative for a particular EI number, additional samples will not be tested unless there is consultation between the Public Health Outbreak Team and the ProvLab.
- The Public Health Outbreak Team will contact the ProvLab if the clinical situation for the outbreak has changed and additional testing needs to be done.

Specimen Transport:

Sites must collect specimens as directed by the Public Health Outbreak Team and make their own arrangement for delivery to the laboratory.

- Follow current Provincial Laboratory for Public Health standards for transporting specimens at [Laboratory Test Directory & Collection Information | Alberta Health Services](#).
- AHS managers and staff can access WHS – Transportation of Dangerous Goods (TDG) modules on My Learning Link for more information regarding safe specimen transport. If staff member does not have access to My Learning Link connect with manager to determine where this learning can be accessed.
- The EI number must be included on each requisition so that specimens receive appropriate testing.
- Rural facilities to transport lab specimens to ProvLab as directed by the Public Health Outbreak Team or by the fastest means possible.

Section 4: Outbreak Definitions & Determining Outbreak Type

Once specimen results are available or residents decline specimen collection, a decision can be made on whether the site has met an outbreak, and the type of outbreak occurring.

Table E: COVID-19 Illness

COVID-19 Illness Case Definition	COVID-19 Illness Outbreak Definition
<p>A person with the virus (SARS-CoV-2) that causes COVID-19 by:</p> <ul style="list-style-type: none"> A positive result on a molecular test (i.e., Nucleic acid amplification test (NAAT's) such as polymerase chain reaction (PCR)), loop-mediated isothermal amplification (LAMP) or rapid molecular test) that is Health Canada approved or approved by the lab accreditation body of the jurisdiction in which the test was performed <p>OR</p> <ul style="list-style-type: none"> A positive result on a Health Canada approved rapid/point-of-care (POC) antigen test in a person with clinical illness (any one or more of the following: new or worsening cough, shortness of breath (SOB), sore throat, loss or altered sense of taste/smell, runny nose/nasal congestion, fever/chills, fatigue (significant and unusual), muscle ache/joint pain, headache, nausea/diarrhea) <p>OR</p> <ul style="list-style-type: none"> Two positive results on a Health Canada approved rapid/POC antigen test completed not less than 24 hours of each other in an asymptomatic person. 	<p>2 or more confirmed COVID-19 cases in residents within a 7-day period, with a common epidemiological link[☆]</p> <p><i>☆Epidemiological link means the cases need to have been in the setting (same site/same unit) during their incubation period or communicable period.</i></p>

Table F: Influenza-Like-Illness (ILI)

ILI Case Definition	ILI Outbreak Definition
<p>Syndromic ILI: Acute onset of respiratory illness which includes cough (new or worsening) and one or more of the following symptom:</p> <ul style="list-style-type: none"> fever shortness of breath sore throat myalgia arthralgia prostration <p>In children under 5 GI symptoms may also be present. In people under 5 or 65 and older fever may not be prominent.</p> <p>Pathogen Specific ILI: Positive for non-influenza, non-COVID-19 pathogen from the Respiratory Pathogen Panel (RPP) - see Table I</p>	<p>2 or more cases of ILI in residents within a 7-day period, with a common epidemiological link[☆]</p> <p>AND</p> <p>No respiratory pathogen identified OR only one case of any respiratory pathogen identified (e.g., Influenza; COVID-19; RSV) OR at least 2 cases of a non-influenza, non-COVID-19 respiratory pathogen</p> <p><i>☆Epidemiological link means the cases need to have been in the setting (same site/same unit) during their incubation period or communicable period</i></p>

Note: ILI outbreaks can be either syndromic or pathogen specific.

- Examples of Syndromic ILI outbreaks include:
 - 2 or more residents who meet the syndromic case definition **OR**
 - 1 resident who meets the syndromic case definition PLUS at least 1 other resident who meets the pathogen-specific case definition **OR**
 - 1 resident positive for influenza or COVID PLUS at least 1 other resident who meets the syndromic case definition
- Examples of Pathogen-specific ILI outbreaks include:
 - 2 or more residents with the same non-influenza, non-COVID pathogen from the RPP

Table G: Influenza Illness

Influenza Illness Case Definition	Influenza Illness Outbreak Definition
<p>A person with clinically compatible signs and symptoms (as outlined above under syndromic ILI) and laboratory confirmation of infection with seasonal influenza virus by:</p> <ul style="list-style-type: none"> detection of influenza virus RNA (e.g., via real-time reverse transcriptase polymerase chain reaction [RT-PCR]) <p>OR</p> <ul style="list-style-type: none"> demonstration of influenza virus antigen in an appropriate clinical specimen (e.g., nasopharyngeal/throat swabs) <p>OR</p> <ul style="list-style-type: none"> significant rise (e.g., fourfold, or greater) in influenza IgG titre between acute and convalescent sera <p>OR</p> <ul style="list-style-type: none"> isolation of influenza virus from an appropriate clinical specimen 	<p>2 or more confirmed influenza cases in residents within a 7-day period, with a common epidemiological link[☆]</p> <p><small>☆Epidemiological link means the cases need to have been in the setting (same site/same unit) during their incubation period or communicable period</small></p>

Table H: Gastrointestinal Illness

Gastrointestinal (GI) Illness Case Definition	GI Illness Outbreak Definition
<p>At least ONE of the following criteria must be met and not be attributed to another cause (e.g., <i>Clostridioides difficile</i> diarrhea, medication, laxatives, diet, or prior medical condition):</p> <ul style="list-style-type: none"> 2 or more episodes of diarrhea (i.e., loose, or watery stools) in a 24-hour period, above what is normally expected for that individual <p>OR</p> <ul style="list-style-type: none"> 2 or more episodes of vomiting in a 24-hour period <p>OR</p> <ul style="list-style-type: none"> 1 or more episodes of vomiting AND diarrhea in a 24-hour period <p>OR</p> <ul style="list-style-type: none"> 1 episode of bloody diarrhea <p>OR</p> <ul style="list-style-type: none"> Laboratory confirmation of a known enteric pathogen AND at least one symptom compatible with a GI infection e.g., nausea, vomiting, diarrhea, abdominal pain, or tenderness 	<p>2 or more cases (with initial onset within one 48-hour period) of GI illness with a common epidemiological link[☆]</p> <p><small>☆Epidemiological link means the cases need to have been in the setting (same site/same unit) during their incubation period or communicable period</small></p>

A **mixed respiratory pathogen outbreak** could result when a combination of lab positive respiratory pathogens/viruses are identified in a facility. Similarly, a **mixed pathogen outbreak** could result when virus(es) causing respiratory and gastrointestinal symptoms are co-circulating in a facility.

The Public Health Outbreak Team will advise which recommendations are applied where there is a mixed pathogen outbreak. The general principle of applying the more protective recommendation will be followed.

Section 5: General Recommendations for Confirmed Outbreaks

The strategies outlined in this section are to be implemented for any sites with a confirmed outbreak, regardless of the outbreak type. **The strategies outlined below would be implemented in addition to any strategies outlined in [Section 3](#).** Additional strategies, specific to the type of outbreak, are discussed in subsequent sections. See [Section 9](#) for GI specific outbreak recommendations.

Note: At the time this guide is written, some zone Outbreak Teams will require that GI Outbreaks utilize the Risk Assessment Worksheet and/or the Risk Assessment Matrix.

The Public Health Outbreak Team may choose to recommend outbreak measures that are not discussed in this guide if the outbreak situation warrants additional measures.

5.1 Infection Prevention and Control Measures

- Maintain all measures discussed in previous section. [See Infection Prevention and Control Measures Section 3.1](#)

5.2 Administrative Measures

- Notify appropriate HCW/staff/departments within the site/facility as indicated by internal protocols (e.g., administration, OHS/WH, IPC, pharmacy). This includes identifying any other professions working in the facility/on the outbreak unit that are not HCW (e.g., construction workers) and consulting with the Public Health Outbreak Team regarding appropriate recommendations for these groups.
- Ensure adequate availability of all supplies through notification of appropriate departments.
- Notify Laundry Services and Distribution Services of the increased need for supplies.
- Ensure that HCW/staff (e.g., nursing, allied health, food services, environmental services, pastoral) are advised of relevant recommendations and work restrictions.
- Ensure that outbreak signage [Attachment 5.1](#) has been posted at the entrance of the facility/unit advising HCW/staff, other professions working in the facility that are not HCW, and visitors of necessary precautions.
- **Once the outbreak is declared, daily reporting of cases to the Public Health Outbreak Team** (and to IPC as per Zone requirement) is required.
 - The facility will use the Facility CDC Outbreak Daily Report Portal (RedCap) ([Attachment 5.2](#)) to report respiratory illness for both residents and symptomatic HCW/staff.
 - The facility will use the data collection tool for GI illness ([Attachment 9.1](#)) or a zone-specific line list, as appropriate to report GI illness for both HCW/staff and residents.
 - Facilities are recommended to follow zone specific outbreak processes for reporting GI illness to the EPH team.
 - The Public Health Outbreak Team will provide direction on how to submit case data and data on case updates (such as hospitalizations and deaths).

- Residents requesting a pass to leave a site/facility may do so if the resident is asymptomatic. Residents are recommended to be advised that if they become symptomatic while away from their site/facility, they are recommended to contact their site/facility to request further direction before returning and seek medical attention if required.
 - Exceptions to this recommendation may apply on a case-by-case basis in consultation with the Public Health Outbreak Team and site leadership. Additional precautions such as enhanced hand hygiene, masking and physical distancing are recommended.
- Inform residents' families/guardians/agent of facility outbreak status.
- The Public Health Outbreak Team uses a variety of strategies to help control and manage the outbreak.
 - Outside of the infection prevention and control measures outlined in [Section 3.1](#) other strategies that will be recommended by the Public Health Outbreak Team include:
 - Continuous masking for all HCW/staff
 - Continuous eye protection for all HCW/staff
 - Masking for all visitors and DFSPs
 - Cancelling high risk activities e.g., singing, bus outing and large group activities
 - Residents to wear masks and practice physical distancing during low-risk activities (e.g., playing cards, participating in an activity or craft) during the outbreak.
 - Additional measures maybe recommended at the discretion of the Public Health Outbreak team including but not limited to:
 - Physical distancing in communal dining area
 - HCW/staff active screening for symptoms prior to each shift during the outbreak
 - Active screening of visitors and DFSPs prior to entering the facility or visiting residents during the outbreak
 - Resident health screening upon return from absence
 - Masking for unvaccinated residents upon return from absence
 - Quarantine and or active screening for residents' admissions upon return from other health settings if that other site is on outbreak.
 - During a COVID-19 outbreak close contact identification and management of residents or HCW/staff may be implemented to control the outbreak

5.3 Resident Restrictions

- Maintain all measures discussed in [Section 3.3](#) including isolation of symptomatic residents and additional precautions.
- Symptomatic residents will be allowed to attend medically necessary activities or appointments. They are recommended to wear an appropriate mask (as tolerated). Ensure transport staff and the receiving facility/unit is notified of the outbreak so that appropriate precautions can be taken for the resident on arrival. Virtual visits are recommended to be arranged when possible.
- Symptomatic residents are recommended to avoid contact with other residents.

5.4 Restrictions to Admissions / Transfers / Discharges on Affected Units/Site

- The facility / unit's ability to accept admissions / transfers / discharges (e.g., whether the unit is open or restricted) will be determined by the Public Health Outbreak Team at the time the outbreak is declared.
 - Consult the Public Health Outbreak Team when issues pertaining to admission, discharge and transfers arise during an outbreak.
- If the Public Health Outbreak Team advises that the facility / unit status is "open," then admissions, transfers, and discharges may proceed following usual non-outbreak processes.
- The Public Health Outbreak Team may determine that the facility / unit is "restricted." This means that admissions (including new admissions to the outbreak affected unit/site), transfers (including transfers from the affected unit/site to a different site), and discharges are recommended to be paused or delayed while these restrictions are in place. However, implementation may not be possible or recommended due to resident circumstances or operational need (including bed pressures). If a site with restrictions believes that an admission, transfer, or discharge is recommended to proceed despite restrictions, sites are recommended to complete a [Risk Assessment Worksheet](#) and follow the [Risk Assessment Matrix as appropriate per zone policy](#).
- The scope of unit restrictions is typically dependent on the extent of the outbreak activity within the facility (one unit, one floor, one wing or the entire facility), the ability to cohort HCW/staff to affected areas, and severity of the outbreak (e.g., many residents and HCW/staff affected, new cases continue to develop despite implemented control measures).

Restrictions regarding resident admissions/re-admissions/transfer and activities are ONLY modified or lifted by the Public Health Outbreak Team. In the event that restriction to admissions/transfers is unduly impacting the availability of acute care beds for individuals requiring urgent care, or because of the expressed informed individual resident or family choice in keeping with AHS's commitment to People Centered Care, the Public Health Outbreak Team will assess the circumstances surrounding the restriction, including the degree of risk to the full spectrum of individuals requiring care.

5.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site

- For residents who were hospitalized prior to the outbreak, or who were hospitalized during an outbreak for an unrelated condition (e.g., fracture), their ability to return to the site will depend on the status of any restrictions applied to the DSL/LTC/Hospice site. See [Section 5.4](#) above, for more details.
 - If the outbreak DSL/LTC site is "open," then the resident can return to their site
 - If the outbreak DSL/LTC site is "restricted," sites are recommended to complete a [Risk Assessment Worksheet](#) and follow the [Risk Assessment Matrix](#)
- If a resident was hospitalized due to illness from the outbreak pathogen, they may return to their home site immediately upon discharge. The need for isolation is recommended to be assessed based on the resident status at the time of return to the site.

5.6 Transfers from an Outbreak LTC, DSL or Hospice site to an Acute Care Site

- If a resident requires acute medical attention or treatment off site (e.g., emergency room, urgent care, dialysis), the outbreak LTC, DSL or Hospice site is recommended to notify the EMS Dispatcher or the transport staff, and the receiving care facility that the resident is being transferred from a LTC site experiencing an outbreak. The transport staff and the facility receiving the resident can then ensure the appropriate precautions are in place when the resident arrives at the hospital/treatment centre. If tolerated, symptomatic residents are recommended to wear an appropriate mask during transfer.

5.7 Group/Social Activities and Non-Resident Events

- Symptomatic residents are not recommended to participate in group or social activities with other residents.
 - Symptomatic residents may continue to participate in social activities with visitors and DFSPs. This includes going outdoors and participating in 1:1 activities as long as additional precautions are followed such as enhanced hand hygiene, masking and physical distancing.
- Consult the Public Health Outbreak Team for recommendations on whether routine group activities may continue for asymptomatic residents.
- For respiratory illness outbreaks, outbreak measures (such as physical distancing, masking, hand hygiene, enhanced surveillance, etc.) may be used for low-risk group activities (e.g., art class, bingo, card games). These activities may continue at the discretion of the facility in consultation with the Public Health Outbreak Team.
- High risk group activities (e.g., singing, bus outing and large group activities) are recommended to be postponed.
- During GI outbreaks, all group activities are recommended to be postponed ([Section 9.7](#)).
- Essential medical treatment activities including but not limited to rehabilitation, physical or group therapy are recommended to be facilitated by the site whenever possible with appropriate precautions in place.
- It is recommended that non-resident events previously booked for areas in the outbreak DSL/LTC/Hospice site (e.g., meetings) be cancelled/postponed.

5.8 Nourishment Areas/Sharing of Food

- Symptomatic residents are recommended to have meals in their rooms.
- The Public Health Outbreak Team will provide direction on any modifications, which may include:
 - Close buffet lines, or, have staff dispense foods from the buffet onto plates for residents.
 - Cease Family-style meal service.
 - Pre-set the tables in common dining areas to minimize resident handling of multiple sets of cutlery.
 - Remove shared food containers from dining areas (e.g., shared pitchers of water, shared coffee cream dispensers, salt & pepper shakers, etc.).
 - For snack programs, dispense snacks directly to residents and use pre-packaged snacks only.
 - If using single service packets of condiments, provide packet directly to each resident, rather than self-serve in a bulk container.
 - Cease activities involving resident participation in food preparation.

- Use of disposable plates and cutlery by symptomatic residents is not required.
- Close the kitchen/nourishment areas accessed by residents/visitors and ensure there is no communal sharing of food in outbreak areas.

5.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3)

- If the outbreak facility operates an Adult Day Program, discuss continuance or stopping of this activity with the OMT/Public Health Outbreak Team.
- Generally, the Public Health Outbreak Team will recommend that Adult Day Programs continue to operate in a facility with an ongoing outbreak if:
 - The Adult Day Program is operating in an area physically separate from areas of the facility in which there have been resident cases with symptoms.
 - Residents attending the Adult Day Program do not socialize with the residents from the outbreak facility.
 - Adult Day Program HCW/staff do not provide care in the areas of the facility in which there have been outbreak cases.

5.10 Visitors and Designated Family/Support Person(s)

- Post outbreak signage [Attachment 5.1](#) at the entrance(s) of the facility/unit advising HCW/staff and visitors of necessary precautions.
- Encourage visitors to postpone visiting if possible. Visitors who choose to visit are recommended to be advised of potential risk of exposure to practice good hand hygiene and limit visit to one (1) resident and exit the facility immediately after the visit.
- Visitation restrictions may be recommended by the Public Health Outbreak Team, depending upon the type of Outbreak confirmed (see additional details in the syndrome or pathogen specific sections that follow).
- A facility/unit may make an operational decision to implement visitation restrictions even if visitation restrictions are not recommended by the Public Health Outbreak Team. However, the facility/unit is recommended to consult with the Public Health Outbreak Team prior to implementing this operational decision and are recommended to consider the impact of the decision on resident and family well-being.
- If visitation restrictions are recommended by the Public Health Outbreak Team, exceptions may still be considered under certain circumstances (e.g., if a resident is receiving end of life care).
- [Refer to Section 3.10 - Visitors and Designated Family/Support Persons](#)

5.11 Volunteers

- Advise volunteers of the potential risk of acquiring illness during outbreaks.
- Have volunteers who continue to assist during an outbreak follow the same control measures as HCW/staff (Section 5.12 below).

5.12 HCW/Staff Outbreak Measures (including volunteers, students, physicians)

- See [Section 3.12](#).
- It is recommended that HCW/staff who work in more than one facility/unit inform any additional facilities or units where they work that an outbreak is in progress to determine whether they are recommended to continue to work in all settings.
- HCW/staff are recommended to monitor themselves for signs and symptoms of illness.

- Symptomatic HCW/staff are recommended to take an at-home rapid antigen test. See [Table D](#). Specific recommendations for unimmunized HCW/staff work restrictions during a confirmed influenza outbreak are found in [Section 8](#).
- Specific recommendations for HCW/staff work restrictions during a GI outbreak are found in [Section 9](#).

5.13 Specimen Collection

- Respiratory illness: symptomatic residents are recommended to have a NP/throat swab collected to determine the causative agent.
 - The number of residents and type of tests that are required for outbreak management will be determined in consultation with the Public Health Outbreak Team.
- GI illness: Sites must collect specimens as directed by the Public Health Outbreak Team and arrange for delivery of specimens to the laboratory. It is valuable to collect stool specimens from cases during outbreaks to try and identify the etiology, if possible. Typical GI illness outbreak stool specimen collection procedures are found in [Attachment 3.1](#).
- Notify the Public Health Outbreak Team if:
 - If there is new symptom presentation among residents within the outbreak site
 - If the outbreak extends beyond original site/unit

5.14 Enhanced Environmental Cleaning and Disinfection

- See [Section 3.14](#)

Roles and Responsibilities for Confirmed Outbreaks

Note: Due to the complex nature of LTC/DSL/Hospice settings, staffing and resident populations, the individual fulfilling the roles and responsibilities within a site may vary from what is outlined below (e.g., some sites may combine the roles of IPC and WHS, or may have designated staff to fulfill these roles)

The Public Health Outbreak Team (MOH, CDC, EPH)

- Provides direction on outbreak management, including infection prevention and control measures, management of residents, management of HCW/staff, and restrictions.
- Provides direction on the need for specimen collection (if applicable)
- Reinforces the need to maintain outbreak control measures and infection control practices as outlined in [Section 3](#).
- Sends out outbreak notifications and alerts as appropriate; and if relevant posts provincial and national Public Health alerts on Canadian Network for Public Health Intelligence (CNPHI).
- Reports outbreak to Alberta Health (AH), and to AHS Senior Public Health Executive (as established within Zone protocol).
- Reviews daily outbreak data received from outbreak site, monitors outbreak progress and provides consultation to the outbreak site when appropriate. The Public Health Outbreak Team is recommended to share timely outbreak status reports with Acute Care, Non-Designated Supportive Living, DSL, and/or LTC partners in their Zone (including AHS and contracted operators), as appropriate. It must be ensured that any shared reports comply with AHS corporate policies for collection, access, use, retention, and disclosure of personal and health information under the care and control of AHS.

- Responds to media inquiries in consultation with AHS Communications Media Advisor.
- Tracks all outbreak samples.
- Participates in OMT meetings.

Infection Prevention and Control (Infection Control Professional and/or Infection Control Designate)

Note: *In the absence of formal IPC or site ICD coverage, facility administration/manager designates responsibility for these roles. This role may be fulfilled by a single individual at the site, or by a combination of a site Infection Control Designate and supporting groups such as organizational/zone/AHS ICP.*

- Directs the implementation of additional outbreak control strategies as indicated according to the type and scope of the outbreak, including prophylaxis if recommended by the MOH, in consultation with site Medical Lead.
- Notifies appropriate HCW/staff within the facility of outbreak as indicated by internal protocols (e.g., administration, medical director, pharmacy, etc.).
- Obtains reports on the clinical status of all affected individuals and works with HCW/staff to identify new cases.
- Provides the Public Health Outbreak Team with status updates of outbreaks within their facility, including relevant immunization status of residents.
- Completes or delegates daily submission of accurate and updated illness data related to the outbreak. The Public Health Outbreak Team will advise IPC, at the time of the outbreak, how daily illness data will be submitted to the Public Health Outbreak Team. See below:
 - Facility CDC Outbreak Daily Report Portal (RedCap) [Attachment 5.2](#)
 - Data collection for GI illness [Attachment 9.1](#)
- Participates in OMT meetings.
- Follows the Public Health Outbreak Team direction regarding the need for additional or subsequent specimens.

Facility Administration/Facility Management or their Designate

- Responsible for maintaining operations to provide optimal care and services for residents during an outbreak.
- Collaborates with IPC/ICD and the Public Health Outbreak Team to implement outbreak management strategies.
- Works collaboratively with the Public Health Outbreak Team to disseminate information to HCW/staff, residents, students, other individuals working in the facility that are not HCW/staff, other departments, and families, as needed.
- Communicates with HCW/staff, residents, families, volunteers, students, and visitors within the facility as appropriate.
- Consult OHS/WHs and/or IPC/ICD as needed.
- Participates on OMT as appropriate.
- Consults with the Public Health Outbreak Team on issues pertaining to admission, discharge, and transfers during an outbreak.
- Notifies senior management within the site as indicated by internal protocols.
- Communicates to appropriate stakeholders outside of the facility.
- Complies with unit/facility restrictions as recommended by the Public Health Outbreak Team.
- Ensures outbreak control strategies are maintained until the outbreak is declared over.

Site/Unit Manager/Designate

- Is the single point of contact for HCW/staff calling in/reporting illness/symptoms.
- Anticipates and provides adequate resources for outbreak management.
- Ensures proper collection of additional or subsequent specimens as directed by the Public Health Outbreak Team and makes own internal arrangements for transporting specimens to the lab.
- Communicates outbreak status to other programs that may be impacted by the Outbreak (i.e., Adult Day Programs, Childcare programs or any other programs that operate within the facility that might be impacted by the outbreak).
- Participates on OMT as appropriate.

Occupational Health/Workplace Health and Safety/Designate

Note: *In the absence of formal AHS WHS or site OHS coverage, facility administration/manager designates responsibility for these roles.*

- Maintains close communication with Frontline Unit/Site Manager with respect to HCW/staff work restrictions/return to work assessments.
- Assesses HCW/staff suitability for return to work.
- Provides information to HCW/staff about work restrictions. The specific restrictions will be based upon the type of outbreak – see pathogen and syndrome specific sections which follow.
- Follows recommendations from OMT, as directed by the Public Health Outbreak Team and/or WHS/OHS.
- Supports illness assessment and surveillance of HCW/staff from outbreak unit as required.
- Identifies HCW/staff who may be at risk of exposure and infection (e.g., unimmunized).
- Participates in OMT meetings.
- Assesses for HCW/staff PPE breaches or hand hygiene breaches as required.

Onsite HCW/Staff (hired directly by the site; contracted HCW/staff, AHS HCW/staff)

- Works collaboratively to ensure ongoing monitoring and surveillance at the site to identify newly symptomatic illness in residents according to [Table A](#) and [Table B](#)
- Facilitates implementing outbreak control strategies as indicated in this document.
- Coordinates the collection of clinical specimens as appropriate, under the direction of the Public Health Outbreak Team.
- May participate in OMT meetings.
- Works collaboratively to ensure ongoing monitoring and surveillance at the site to identify new symptomatic residents.
- Works collaboratively to provide the Public Health Outbreak Team with daily status updates on the outbreak.
- Maintains clear and consistent channels of communication within the outbreak site as per established Zone practices.
- Liaises with Most Responsible Health Practitioner, as necessary.
- Provides services to care and treat residents in place as per [Infection Prevention and Control Risk Assessment \(IPCRA\)](#).
- Authorizes additional professional and support services, as needed, to manage the outbreak. This may include such things as: assessment/monitoring of symptomatic residents, laundry, bathing, meal tray delivery and environmental cleaning.

- Communicates outbreak status to other programs that may be impacted by the Outbreak (i.e., Adult Day Programs, Childcare programs or any other programs that operate within the facility that might be impacted by the outbreak).
- Reports PPE breaches or hand hygiene breaches to supervisor or Occupational Health/ Workplace Health and Safety as appropriate.



Provincial Laboratory for Public Health (ProvLab)

- Ensures the Public Health Outbreak Team and IPC/ICD (if noted on the requisition) receive timely results of outbreak specimens.
- Tracks all outbreak samples.

Attachment 5.1: Outbreak Signage

Sites may print black and white or colour outbreak signage from:

<https://www.albertahealthservices.ca/assets/healthinfo/ipc/if-hp-ipc-facility-outbreak-stoplight-poster-colour.pdf>



Attachment 5.2: Facility CDC Outbreak Daily Report Portal (RedCap)

Site Under Investigation / Confirmed Outbreak

Facility CDC Outbreak Daily Report Portal (RedCap) Email Template

Hello,

As discussed, [Site Name] is under investigation for a potential outbreak / has a confirmed outbreak of [pathogen]. You are required to submit daily notification, by 10:00 AM, of the following to the Public Health Outbreak Team for reporting purposes:

- Report if no new cases in staff or residents in the past 24 hours
- Newly symptomatic residents
- Newly symptomatic staff (includes contracted staff)
- Newly positive lab results (including positive Rapid Antigen Tests)
- Report new hospitalizations or deaths due to the illness in residents or staff, including those previously reported as only symptomatic
 - Include any death that occurs within 30 days of the positive lab
 - Include any death that occurs greater than 30 days from the positive lab AND COVID is attributed as a primary or secondary cause of death

The submission is completed electronically through the online portal at:

<https://redcap.link/FacilityCDCOutbreakReport2022>

You must enter the EI number for your site with each entry and provide it when you call, so be sure to have this information with you. Your EI is: [202X-EI-XXXXX]

The portal will require the following information:

- Demographic information name, DOB, ULI/PHN for resident and name; DOB, phone number for staff.
- Onset date of illness, symptoms.
- Whether a swab has been obtained, and date of swab if obtained.
- Whether individual is hospitalized; and
- If an individual has died (include date of death).
- To report lab confirmed cases in residents or staff (including asymptomatic cases):
 - Begin entry as if symptomatic by choosing "Newly Symptomatic resident/staff"
 - Use specimen collection date as onset date
 - Select "none" for all symptom lists if asymptomatic

Note: For 'Date of Birth' field, the clickable calendar does not have a drop-down option for years prior to 1920, however years prior to 1920 can be manually added in the DOB field.

Also attached is a letter from the Medical Officer of Health describing the legal authority to release the requested line list information to the Public Health Outbreak Team, for the purposes of outbreak management under the Alberta Public Health Act for your records.

If you have any questions regarding submission of the information through the portal, please email the CDC COVID outbreak email: CDOutbreak@albertahealthservices.ca.

Section 6: Confirmed COVID-19 Outbreak

COVID-19 Outbreak Definition

Outbreak definition: 2 or more confirmed COVID-19 cases in residents (as per [Table A](#)) that:

- Have an epidemiological link between the cases, and
- Occur within a 7-day period.

Duration of Outbreak

- The outbreak remains open for 14 days after the onset of the last resident case (will close on day 15).

The following are baseline or minimum recommendations for a confirmed COVID-19 outbreak. Additional or more restrictive recommendations may be put in place at the discretion of the Public Health Outbreak Team or the facility in consultation with the Public Health Outbreak Team.

6.1 Infection Prevention and Control Measures

- See [Section 3.1](#)
- Once a COVID-19 outbreak is declared, HCW/staff are recommended to use continuous masking and eye protection for the duration of the outbreak. Additional precautions are necessary if performing aerosol-generating medical procedures (AGMP) or a resident is on airborne precautions. [COVID-19 Aerosol-Generating Medical Procedure Guidance Tool | Alberta Health Services](#).
- If resident tests positive for COVID-19 or another organism (see [Table I](#) for isolation recommendations) maintain full IPC precautions until the resident is released from isolation as directed by IPC/ICD or the Public Health Outbreak Team.
- All HCW/staff and residents in outbreak setting are recommended to be made aware of the outbreak and be advised to monitor for symptoms.
- HCW/staff shall wear a surgical/procedure mask, seal-checked KN95 or fit-tested N95 respirator (mask) in accordance with the [Infection Prevention and Control Risk Assessment \(IPCRA\)](#)
- HCW/staff are recommended to report PPE breaches or hand hygiene breaches to supervisor or Occupational Health/ Workplace Health and Safety as appropriate.

6.2 Administrative Measures

- Antiviral treatments may be available for COVID-19. Sites are recommended to have policies in place to ensure residents that meet the criteria have timely access to the medication – see [COVID-19 Outpatient Treatment | Alberta Health Services](#)
- Post signs that facility is on outbreak at entrances [Attachment 5.1](#).
- **Once the outbreak is declared, daily reporting of cases to the Public Health Outbreak Team** (and to IPC as per Zone requirement) is required.
 - The facility will use the Facility CDC Outbreak Daily Report Portal (RedCap) ([Attachment 5.2](#)) to report respiratory illness for both residents and symptomatic HCW/staff.
 - The Public Health Outbreak Team will provide direction on how to submit case data and data on case updates (such as hospitalizations and deaths).

6.3 Resident Restrictions

- Residents who test positive for COVID-19 are recommended to remain in their room with appropriate precautions as per the [Infection Prevention and Control Risk Assessment \(IPCRA\)](#) for a minimum 5 days or until symptoms improve AND they are afebrile for 24 hours without the use of fever reducing medications, **whichever is longer**.
 - A mask must be worn for an additional 5 days when the resident leaves their room.
 - During this 5-day masking period residents must return to their room whenever the mask needs to be removed (e.g., meals and snacks).
 - If a mask is not tolerated, the resident is recommended to complete 10 days of isolation in their room.

6.4 Restrictions on Affected Unit/Site

- Facility / unit status (open or restricted) will be determined by the Public Health Outbreak Team. In some Zones, the Public Health Outbreak Team may permit admissions and transfers under certain conditions, while in other Zones the affected unit/site may be restricted for the duration of the COVID-19 outbreak.
- If the Public Health Outbreak Team advises that the facility/unit status is “open,” then admissions, transfers, and discharges may proceed following usual non-outbreak processes.
- The Public Health Outbreak Team may determine that the facility/unit is “restricted.” This would mean that admissions, transfers, and discharges are recommended to be paused or delayed while these restrictions are in place. However, implementation may not be possible or recommended due to resident circumstances or operational need (including bed pressures). If a site with restrictions believes that an admission, transfer, or discharge is recommended to proceed despite restrictions, sites are recommended to complete a [Risk Assessment Worksheet](#) and follow the [Risk Assessment Matrix](#) as per the zone process.

6.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site

- For residents who were hospitalized prior to the outbreak, their ability to return to the site will depend on the status of any restrictions applied to the DSL/LTC/Hospice site. See [Section 6.4](#) above for more details.
- If a resident was hospitalized due to COVID-19, they may return to their home site immediately upon discharge.
- If a resident was hospitalized during an outbreak for an unrelated condition (e.g., fracture), their ability to return to the site will depend on the status of any restrictions applied to the DSL/LTC/Hospice site. See [Section 6.4](#) above for more details.

6.6 Transfers from an outbreak facility to an acute care site

- See [Section 5.6](#)

6.7 Group / Social Activities and Other Events

- Symptomatic residents are not recommended to participate in group or social activities with other residents.
 - Symptomatic residents may continue to participate in social activities with visitors and DFSPs. This includes going outdoors and participating in 1:1 activities as long as additional precautions are followed such as enhanced hand hygiene, masking and physical distancing.
- Residents that are on isolation are advised to remain in their room but may attend appointment as outlined in [Section 5.3](#).
- Consult the Public Health Outbreak Team for recommendations on whether routine group activities may continue for asymptomatic residents.
- For COVID-19 outbreaks, outbreak measures (such as physical distancing, masking, hand hygiene, enhanced surveillance, etc.) may be used for low-risk group activities (e.g., art class, bingo, card games). These activities may continue at the discretion of the facility in consultation with the Public Health Outbreak Team.
- High risk group activities (e.g., singing, bus outing and large group activities) are recommended to be postponed.
- Essential medical treatment activities including but not limited to rehabilitation, physical or group therapy is recommended to be facilitated by the site whenever possible with appropriate precautions in place.
- It is recommended that non-resident events previously booked for areas in the outbreak LTC/DSL/Hospice site (e.g., meetings) be cancelled/postponed.
- Personal services (e.g., hair styling) are allowed to continue for asymptomatic residents. The personal service provider is required to wear a mask and eye protection and the resident receiving the service is also recommended to wear a mask when possible. PPE will be provided by the facility.
 - Personal services are recommended to be provided to one resident at a time for residents from outbreak units.

6.8 Nourishment Areas / Sharing of Food

- The Public Health Outbreak Team will provide direction on any modifications, which may include moving to single serve items and discontinuing family style dining.
- Group dining may continue – with IPC (such as physical distancing and hand hygiene) measures in place.

6.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3)

- See [Section 5.9](#)

6.10 Visitors and Designated Family/Support Person(s)

- See [Section 3.10](#) and [Section 5.10](#).

6.11 Volunteers

- Advise volunteers of the potential risk of acquiring illness during outbreaks.
- Have volunteers who continue to assist during an outbreak to follow the same control measures as HCW/staff. See [Section 6.12](#)

6.12 HCW/Staff Outbreak Measures (Including volunteers, students, and physicians)

- See [Section 3.12](#) and [Section 5.12](#).
- Encourage all HCW/staff to get all recommended COVID-19 vaccines.
- HCW/staff with a household member (or other close contacts) that is COVID-19 positive is recommended to monitor for symptoms for 7 days after last exposure.
- Once a COVID-19 outbreak is declared, HCW/staff are recommended to use continuous masking and eye protection.
- Additional precautions are necessary if performing aerosol-generating medical procedures (AGMP). [COVID-19 Aerosol-Generating Medical Procedure Guidance Tool | Alberta Health Services](#).

6.13 Specimen Collection

- After the COVID-19 outbreak has been confirmed, additional specimen collection on newly symptomatic residents is not recommended unless directed by the Public Health Outbreak Team
- Notify the Public Health Outbreak Team:
 - If new symptom presentation among residents within the outbreak site
 - If the outbreak extends beyond the original site/unit
- Management of resolved COVID-19 Cases:
 - Previous cases within 90 days are NOT to be retested unless directed by the Public Health Outbreak Team.

6.14 Enhanced Environmental Cleaning and Disinfection

- See [Section 3.14](#)

Roles and Responsibilities for Confirmed Outbreaks

Only actions and tasks above and beyond those discussed in previous sections are listed here. See [Section 5 Roles and Responsibilities for Confirmed Outbreaks](#) for additional details.

Note: Due to the complex nature of LTC/DSL/Hospice settings, staffing and resident populations, the individual fulfilling the roles and responsibilities within a site may vary from what is outlined below (e.g., some sites may combine the roles of IPC and WHS, or may have designated staff to fulfill these roles)

The Public Health Outbreak Team (MOH, CDC, EPH)

- Provides direction on restrictions to admissions / transfers / discharges to the outbreak unit / facility.
- Provides direction on isolation of resident cases.
- Provides direction on management of HCW/staff.
- Provides direction on changes to activities (if applicable) within the unit / facility

Infection Prevention and Control (Infection Control Professional and/or Infection Control Designate)

Note: In the absence of formal IPC or site ICD coverage, facility administration/manager designates responsibility for these roles. This role may be fulfilled by a single individual at the site, or by a combination of a site Infection Control Designate and supporting groups such as organizational/zone/AHS ICP.

- Directs the isolation of resident cases.
- Identifies high risk activities which are recommended to be stopped during the outbreak.

Facility Administration/Facility Management or their Designate

- Supports assessment of resident cases in determining whether site meets outbreak definition.
- Anticipates and manages the impact of HCW/staff exclusion on site operations.
- Ensures that, if recommended, restrictions to admissions / transfer / discharges to the outbreak facility are implemented.
- If facility has restrictions to admissions / transfers / discharges, oversees completion of Risk Assessment Worksheets for operationally necessary resident moves.
- Communicates the status of the facility (open or restricted) to partners and stakeholders.

Site/Unit Manager/Designate

- Provides information about HCW/staff shift patterns (when HCW/staff member last onsite and onsite during infectious period) to the Public Health Outbreak Team and OHS / WHS
- Collects information about HCW/staff immunization status. Shares this information with OHS / WHS and the Public Health Outbreak Team.
- Supports assessment of resident cases in determining whether unit meets outbreak definition.
- Anticipates and manages the impact of HCW/staff exclusion on unit operations.
- Ensures that, if recommended, restrictions to admissions / transfer / discharges to the outbreak unit are implemented.
- If unit has restrictions to admissions / transfers / discharges, oversees completion of Risk Assessment Worksheets for operationally necessary resident moves.
- Communicates the status of the unit (open or restricted) to partners and stakeholders.

Occupational Health/Workplace Health and Safety/Designate

Note: *In the absence of formal AHS WHS or site OHS coverage, facility administration/manager designates responsibility for these roles.*

- Participates in unit/site investigation to determine if HCW/staff cases are recommended to be linked to the site investigation and communicates conclusions to the Public Health Outbreak Team
- Confirms HCW/staff cases and implements HCW/staff work restrictions as required.

Onsite HCW/Staff (hired directly by the site; contracted HCW/staff, AHS HCW/staff)

- Each HCW/staff member is responsible for reporting their personal immunization status with Facility Administration / Facility Management and / or Occupational Health / Workplace Health and Safety.
- Complies with work restrictions, if applicable

Provincial Laboratory for Public Health (ProvLab)

- If applicable, completes additional testing on COVID-19 outbreak specimens to identify variants of interest or variants of concern.

Section 7: Confirmed Influenza-Like Illness Outbreak

7A. Influenza-Like Illness (ILI) Outbreak Definition

- 2 or more cases of ILI in residents within a 7-day period, with a common epidemiological link

AND

No respiratory pathogen identified OR 1 case of any respiratory pathogen identified (e.g., Influenza; COVID-19; RSV) OR at least 2 cases of a non-influenza, non-COVID-19 respiratory pathogen.

7B. Syndromic or Pathogen Specific Influenza-Like Illness (ILI) Outbreaks

- An ILI outbreak can be syndromic (based only on the symptom presentation of resident cases - see [Table F](#)). However, an ILI outbreak may also have a non-influenza, non-COVID respiratory pathogen identified. Examples of common respiratory pathogens can be found in [Table I](#)

7C. Duration of Outbreak

- If a non-influenza, non-COVID-19 respiratory pathogen is identified, the outbreak will remain open for a single incubation period for that pathogen (See [Table I](#) for incubation periods for common respiratory pathogens). For example, the incubation period for RSV is 2 to 8 days, so an RSV outbreak would close the 9th day following the last case onset.
- If no pathogen is identified, an ILI outbreak would remain open for 7 days (would close the 8th day following the last case onset).

Table I: Organisms Commonly Associated with Respiratory Illness

(Reference: [IPC Diseases and Conditions Table for Management of Residents in Acute Care](#), [IPC Diseases and Conditions Table Recommendations for Management of Residents Continuing Care](#) and [Alberta Public Health Disease Management Guidelines - COVID-19](#))

ORGANISM	CLINICAL PRESENTATION /SYMPTOMS	INFECTIOUS SUBSTANCE/ HOW IT IS TRANSMITTED	INCUBATION PERIOD	PERIOD OF COMMUNICABILITY	OUTBREAK RESTRICTIONS/ RECOMMENDATIONS FOR SITES	PRECAUTIONS REQUIRED
INFLUENZA, SEASONAL TYPE A OR B	Sudden onset of fever, cough, muscle aches, fatigue, sore throat, runny nose & sneezing. Note: fever may not be prominent in those >65 years.	Infectious substance is respiratory secretions. Direct contact, indirect contact, and large droplets - Person to person by large droplets or direct contact with articles recently contaminated with respiratory secretions.	1-3 days**	For duration of symptoms – usually 3-7 days from onset of symptoms.	Residents who test positive are recommended to remain on isolation for 5 days from the onset of acute illness OR until they are over their acute illness and have been afebrile for 24 hours, whichever is longer. Refer to Section 8: Confirmed Influenza Outbreak for further information.	Contact and Droplet
COVID-19 (SARS-CoV-2)	Fever, new onset or worsening of cough, new or worsening shortness of breath or difficulty breathing, sore throat, runny nose, and loss of sense of smell or taste. Extended symptoms include chills, painful swallowing, stuffy nose, headache, muscle, or joint ache, feeling unwell, fatigue or severe exhaustion, nausea, vomiting, diarrhea or unexpected loss of appetite, conjunctivitis.	Infectious substance is respiratory secretions. Droplet, direct and indirect contact. Person to person by large droplets or smaller droplets (aerosols) that in certain circumstances (such as with AGMPs, indoor spaces with poor ventilation) have the potential to be infectious over longer distances; or direct contact with articles recently contaminated with respiratory secretions.	1-14 days	May begin up to 48 hours prior to symptom onset and continue throughout the symptomatic period. Current studies indicate that by day eight of illness/symptoms, no live virus was recovered from patients with upper respiratory tract disease or limited lower respiratory tract disease. People with more severe disease are likely to be infectious for a few days longer.	Residents who test positive for COVID-19 must isolate in their room with appropriate precautions as per the Infection Prevention and Control Risk Assessment (IPCRA) for a minimum 5 days or until symptoms improve AND they are afebrile for 24 hours without the use of fever reducing medications, whichever is longer . A mask must be worn for an additional 5 days when the resident leaves their room. Refer to: Section 6: Confirmed COVID-19 Outbreaks for further information.	Modified Respiratory Precautions

ORGANISM	CLINICAL PRESENTATION /SYMPTOMS	INFECTIOUS SUBSTANCE/ HOW IT IS TRANSMITTED	INCUBATION PERIOD	PERIOD OF COMMUNICABILITY	OUTBREAK RESTRICTIONS/ RECOMMENDATIONS FOR SITES	PRECAUTIONS REQUIRED
RESPIRATORY SYNCYTIAL VIRUS (RSV)	Runny nose, coughing, sneezing, fever, wheezing.	Infectious substance is respiratory secretions. Direct contact, indirect contact, and large droplets - Person to person by large droplets or direct contact with articles recently contaminated with respiratory secretions.	2 to 8 days	For duration of symptoms.	Confirmed or symptomatic cases are recommended to remain in their rooms for the duration of the illness , which is the resolution of acute respiratory infection symptoms or return to baseline. Admission/transfer restrictions only when recommended by local MOH.	Contact and Droplet
PARAINFLUENZA Type 1, 2, 3, 4	Fever, runny nose, cough, sneezing, wheezing, sore throat, croup, bronchitis.	Infectious substance is respiratory secretions. Direct contact, indirect contact, and large droplets - Person to person by large droplets or direct contact with articles recently contaminated with respiratory secretions.	2 to 6 days	1-3 weeks	Confirmed or symptomatic cases are recommended to remain on precautions for 5 days from the onset of acute illness OR until they are over the acute illness and have been afebrile x 48 hours (whichever is longer) Admission/transfer restrictions only when recommended by local MOH.	Contact and Droplet
HUMAN METAPNEUMO-VIRUS (hMPV)	Cough, fever, nasal congestion, shortness of breath.	Infectious substance is respiratory secretions Direct contact, indirect contact and large droplets - Person to person by large droplets or direct contact with articles recently contaminated with respiratory secretions.	3 to 5 days	For duration of symptoms	Confirmed or symptomatic cases are recommended to remain in their rooms for the duration of the illness , which is resolution of acute respiratory infection symptoms or return to baseline. Admission/transfer restrictions only when recommended by local MOH.	Contact and Droplet

ORGANISM	CLINICAL PRESENTATION /SYMPTOMS	INFECTIOUS SUBSTANCE/ HOW IT IS TRANSMITTED	INCUBATION PERIOD	PERIOD OF COMMUNICABILITY	OUTBREAK RESTRICTIONS/ RECOMMENDATIONS FOR SITES	PRECAUTIONS REQUIRED
OTHER COMMON RESPIRATORY VIRUSES SUCH AS: ENTERO/RHINOVIRUS NON-COVID-19 CORONAVIRUSES ADENOVIRUS	Sore throat, runny nose, coughing, sneezing.	Infectious substance is respiratory secretions Direct contact, indirect contact, and large droplets - Person to person by large droplets or direct contact with articles recently contaminated with respiratory secretions	Enteroviruses: usually 2-3 days Non-COVID-19 Coronaviruses: usually 2-4 days Adenovirus: 2-14 days	For duration of symptoms	Confirmed or symptomatic cases are recommended to remain in their room for the duration of the illness , which is resolution of acute respiratory infection symptoms or return to baseline. Admission/transfer restrictions only when recommended by local MOH.	Contact and Droplet

**First day is designated as Day 0; after the first 24 hours is Day 1. The Public Health Outbreak Team declares the facility outbreak to be over 7 days after onset of symptoms in the last case (e.g., closes outbreak on the morning of day 8).

7.1 Infection Prevention and Control

- See [Section 5.1](#).

7.2 Administrative Measures

- After the ILI outbreak has been confirmed, the Public Health Outbreak Team will provide direction on how to report additional resident cases.
- **Once the outbreak is declared, daily reporting of cases to the Public Health Outbreak Team** (and to IPC as per Zone requirement) is required.
 - The facility will use the Facility CDC Outbreak Daily Report Portal (RedCap) ([Attachment 5.2](#)) to report respiratory illness for both residents and symptomatic HCW/staff.
 - The Public Health Outbreak Team will provide direction on how to submit case data and data on case updates (such as hospitalizations and deaths).

7.3 Resident Restrictions

- See [Section 3.3](#)
- Symptomatic residents are recommended to be placed on appropriate precautions. The duration of the precautions depends on the cause of the ILI outbreak.
 - If a specific pathogen has been identified, maintain [Contact and Droplet Precautions](#) for the duration discussed in [Table I: Organisms Commonly Associated with Respiratory Illness](#).
 - If no specific pathogen has been identified, symptomatic residents are recommended to remain in their rooms for the duration of the illness, which is resolution of acute respiratory infection symptoms or return to baseline.
- Symptomatic residents will be allowed to attend medically necessary activities or appointments, they are recommended to wear an appropriate mask (as tolerated). Ensure receiving facility/unit is notified of the potential outbreak so that appropriate precautions can be taken for the resident on arrival. Virtual visits are recommended to be arranged when possible.
- If symptomatic residents cannot remain in their room, they are recommended to avoid contact with other residents in common areas as much as possible.

7.4 Restrictions to Admissions / Transfers / Discharges on Affected Unit / Site

- See [Section 5.4](#).
- In general, facility/unit would remain “open” during an ILI outbreak, meaning that admissions, transfers, and discharges may proceed following usual non-outbreak processes.
 - Note: for mixed pathogens including ILI + Influenza, or ILI + GI the facility/unit would generally be closed.

7.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site

- Unless the Public Health Outbreak Team advises the Unit / Site that is “restricted” (see [Section 7.4](#)), admissions / transfers from acute care may proceed.

7.6 Transfers from an Outbreak Facility to an Acute Care Site

- If a resident requires acute medical attention or treatment off site (e.g., emergency room, urgent care, dialysis), the outbreak facility must notify the EMS Dispatcher, the transport staff (EMS crew) and the receiving care facility that the resident is being transferred from a facility experiencing an ILI outbreak. The transport staff (EMS crew) and the facility receiving the resident can then ensure contact and droplet precautions are in place when the resident arrives at the hospital/treatment centre. If tolerated, symptomatic residents are recommended to wear an appropriate mask during transfer.

7.7 Group / Social Activities and Other Events

- See [Section 5.7](#).
- Unless otherwise directed by the Public Health Outbreak Team, routine group activities may continue for asymptomatic residents.

7.8 Nourishment Areas / Sharing of Food

- See [Section 5.8](#).
- The Public Health Outbreak Team will provide direction on any modifications, which may include moving to single serve items and discontinuing family style dining.
- Group dining may continue.

7.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3)

- See [Section 5.9](#)

7.10 Visitors and Designated Support Person(s)

- See [Section 3.10](#)
- See [Section 5.10](#)

7.11 Volunteers

- Advise volunteers of the potential risk of acquiring illness during outbreaks.
- Have volunteers who continue to assist during an outbreak to follow the same control measures as HCW/staff. See [Section 7.12](#).

7.12 HCW/Staff Outbreak Measures (Including volunteers, students, physicians)

- See [Section 3.12](#) and [Section 5.12](#).

7.13 Specimen Collection

- After the Influenza-Like Illness outbreak has been confirmed, additional specimen collection on newly symptomatic residents is recommended. Consult the Public Health Outbreak Team regarding which specimens and testing is recommended.
- Notify the Public Health Outbreak Team if:
 - If new symptom presentation among residents within the outbreak site
 - If the outbreak moves extends beyond original site

7.14 Enhanced Environmental Cleaning and Disinfection

- See [Section 3.14](#)

Roles and Responsibilities for Confirmed Outbreaks

Roles and responsibilities for an RI outbreak are similar to those in previous sections. See [Section 5 Roles and Responsibilities for Confirmed Outbreaks](#) for additional details.

Section 8: Confirmed Influenza Outbreak

The symptoms of influenza disease are similar to the symptoms for many other respiratory illnesses. See the Health Canada Guidance for further information for identification of suspect influenza cases and indications for early treatment with antivirals: [Flu \(influenza\): For health professionals - Canada.ca](https://www.canada.ca/en/health-canada/services/influenza/flu-influenza-for-health-professionals.html)

Influenza Outbreak Definition

- 2 or more cases of laboratory confirmed influenza in residents, with a common epidemiological link[☆], within a seven-day period

[☆]*Epidemiological link means the cases need to have been in the setting (same site/same unit) during their incubation period or communicable period.*

Duration of Outbreak

- Outbreak remains open for 7 days after symptom onset of most recent influenza case.
- Additional information about influenza disease, including the incubation period and, period of communicability, can be found in [Table I: Organisms Commonly Associated with RI](#)

8.1 Infection Prevention and Control

- See [Section 3.1](#)
- See [Section 3.14](#)

The Outbreak measures listed below are specific to a confirmed influenza outbreak. See [Section 5](#) for additional General Measures recommended during any confirmed outbreak.

8.2 Administrative Measures

- The major interventions for an influenza outbreak include:
 - Antiviral prophylaxis for all residents [Section 8.3](#)
 - Antiviral prophylaxis for unimmunized HCW/staff [Section 8.12](#)
 - Restrictions to admissions, transfers, and discharges [Section 8.4](#)
- **Once the outbreak is declared, daily reporting of cases to the Public Health Outbreak Team** (and to IPC as per Zone requirement) is required.
 - The facility will use the Facility CDC Outbreak Daily Report Portal (RedCap) ([Attachment 5.2](#)) to report respiratory illness for both residents and symptomatic HCW/staff.
 - The Public Health Outbreak Team will provide direction on how to submit case data and data on case updates (such as hospitalizations and deaths).

8.3 Resident Restrictions

- Resident cases and symptomatic residents are recommended to remain on isolation for 5 days from the onset of acute illness OR until they are over their acute illness and have been afebrile for 24 hours, whichever is longer.
- Resident influenza cases and symptomatic residents who become ill during a confirmed influenza outbreak are recommended to receive Oseltamivir (Tamiflu) treatment. See [Appendix A](#) to review Antiviral (Oseltamivir) Dosing Recommendations.

- Asymptomatic residents, regardless of immunization status, are recommended to receive Oseltamivir (Tamiflu) prophylaxis. Antiviral prophylaxis is continued for 7 days after onset of symptoms of the last resident case, usually a minimum of 10 days. See [Appendix A](#) to review Antiviral (Oseltamivir) Dosing Recommendations.
 - In the situation of a mixed outbreak: COVID-19 and influenza, contact the Public Health Outbreak Team to discuss the length of Tamiflu prophylaxis.

8.4 Restrictions on Affected Units/Site

- The Public Health Outbreak Team will make a best practice recommendation that the facility / unit is “restricted.” This means that admissions, transfers, and discharges are recommended to be paused or delayed during the influenza outbreak. However, implementation of this best practice recommendation may not be possible or recommended due to resident circumstances or operational need (including bed pressures). If a site with restrictions believes that an admission, transfer, or discharge is recommended to proceed despite restrictions, sites are recommended to follow the zone Risk Assessment Process and follow the [Risk Assessment Matrix](#)
- For confirmed influenza outbreaks, **admission restrictions will remain in place at minimum for 7 days following the onset of symptoms in the last case**, based on [recommendations from the Association of Medical Microbiology and Infectious Disease \(AMMI\) Canada](#), and as directed by the Public Health Outbreak Team.

Restrictions regarding resident admissions/re-admissions/transfer and activities are **ONLY** modified or lifted by the Public Health Outbreak Team.

If restriction to admissions/transfers is unduly impacting the availability of acute care beds for individuals requiring urgent care, or because of the expressed informed individual resident or family choice (in keeping with AHS’s commitment to People Centered Care), the Public Health Outbreak Team will assess the circumstances surrounding the restriction, including the degree of risk to the full spectrum of individuals requiring care.

8.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site

- If a resident was hospitalized prior to the outbreak or due to an unrelated condition (e.g., fracture), the acute care site and the outbreak site must collaborate on the Risk Assessment Process to explore if the resident may return to the facility. Oseltamivir (Tamiflu) prophylaxis for the resident may be required.
- If a resident was hospitalized due to influenza, they may return to their home site immediately upon discharge.

If an admission/transfer to a LTC, DSL or Hospice site must occur during a confirmed influenza outbreak at the site, following the assessment of the circumstances and consultation with the Public Health Outbreak Team (as described in the box above), LTC, DSL or Hospice HCW/staff are recommended to collaborate with the acute care contact before the resident is discharged. See [Risk Assessment Worksheet](#).

The resident is not recommended to be transferred until the LTC, DSL or Hospice site HCW/staff ensures that:

- The resident/guardian has information on risks associated with the outbreak and consents to the transfer AND
- The resident is immunized AND
- The resident/guardian is able to and agreeable to take antiviral medication as indicated.

8.6 Transfers from an Outbreak Facility to an Acute Care Site

- See [Section 5.6](#)

8.7 Group / Social Activities and Other Events

- See [Section 5.7](#)

8.8 Nourishment Areas / Sharing of Food

- See [Section 5.8](#)

8.9 Operating Adult Day Programs during an Outbreak

- See [Section 5.9](#)

8.10 Visitors and Designated Support Person(s)

- Visitors are strongly encouraged to receive annual immunization for influenza when available.
- Advise those who choose to visit during an outbreak to practice good hand hygiene, visit one (1) resident only and exit the facility immediately after the visit.
- See [Section 5.10](#)
- See [Section 3.10](#)

8.11 Volunteers

- Volunteers are strongly encouraged to receive annual influenza immunization when available.
- Have volunteers who continue to assist during an outbreak follow the same control measures as HCW/staff. See [Section 8.12](#).

8.12 HCW/Staff Outbreak Measures (Including volunteers, students, physicians)

- See [Section 3.12](#) and [Section 5.12](#).
- HCW/staff are strongly encouraged to receive an annual dose of seasonal influenza vaccine when available.

- The Public Health Outbreak Team will advise whether the outbreak influenza strain is covered in the seasonal influenza vaccine. If the outbreak strain is not covered in the seasonal vaccine, the Public Health Outbreak Team may provide additional direction beyond what is described below.

Management of Asymptomatic HCW/Staff who are not immunized.

Recommendations for post-exposure immunization, prophylaxis and/or work restrictions to control influenza A or B outbreaks will be directed by the Public Health Outbreak Team. Oseltamivir (Tamiflu) antiviral treatment and prophylaxis is administered as per the most current Alberta Health Influenza Antiviral Drug Policy. It is the responsibility of the facility to clearly communicate instructions to their HCW/staff on how to access antiviral prophylaxis. See [Appendix A](#) for further information.

Asymptomatic HCW/staff on the outbreak unit/facility who have not received a dose of the current season's vaccine (are considered unimmunized) generally fall into three categories, each subject to work restrictions:

- HCW/staff who are not immunized and who are asymptomatic at time of assessment **and** who agree to be immunized, **but** do not take prophylaxis are recommended to be:
 - Excluded from work for 3 days from the last day of work on the outbreak unit/site.
 - If they remain asymptomatic after 3 days and receive immunization, they may be reassigned to a non-outbreak unit/site for the duration of the outbreak or for 14 days from date of immunization, whichever occurs first.
 - If reassignment of work is not possible, then the HCW/staff is recommended to be excluded from work for 14 days from the time of immunization or for the duration of the outbreak, whichever occurs first.
- **Asymptomatic HCW/staff who are not immunized and do not take the recommended antiviral prophylaxis are recommended to be:**
 - Excluded from working in the affected facility/unit(s) until the outbreak is over,

OR

 - Relocated to a non-outbreak unit/site if they remain asymptomatic after waiting 3 days from the last day of work on the outbreak unit and are not recommended to return to the outbreak unit/site for the duration of the outbreak.
- **Asymptomatic HCW/staff immunized less than 14 days prior to the outbreak and do not take recommended antiviral prophylaxis are recommended to be:**
 - Excluded from working in the affected unit/facility until 14 days from date of immunization, or for the duration of the outbreak whichever occurs first,

OR

 - Excluded from working at any site for 3 working days from the last day of work on the outbreak unit. If they remain asymptomatic after waiting the 3 working days, they can be relocated to a non-outbreak unit until 14 days from the date of immunization or for the duration of the outbreak at the manager's discretion.
- If required, the Public Health Outbreak Team may supply letters that the unit/site can provide to HCW/staff who are not immunized to reinforce their options.
 - [Attachment 8.1](#) MOH Notice to all Unimmunized HCW - Exclusion from Work (SAMPLE) Letter used for Confirmed Influenza Outbreaks
 - [Attachment 8.2](#) MOH Notice to HCW Immunized less than 14 days prior to declaration of Outbreak - Exclusion from Work (SAMPLE) Letter used for Confirmed Influenza
- For asymptomatic HCW/staff, no waiting period is required between starting Oseltamivir (Tamiflu) prophylaxis and returning to work.

- Oseltamivir (Tamiflu) prophylaxis for HCW/staff is NOT publicly funded. Sites are responsible for developing a site policy on who is responsible for the cost of HCW/staff prophylaxis.

Management of Symptomatic HCW/Staff

- If HCW/staff develop symptoms, they are recommended to stay home and follow recommendations in [Section 3.12](#) and [Section 5.12](#).

8.13 Specimen Collection

- After the Influenza outbreak has been confirmed, additional outbreak specimen collection of newly symptomatic residents is not recommended unless directed by the Public Health Outbreak Team
- Notify the Public Health Outbreak Team:
 - If new symptom presentation among residents within the outbreak site
 - If the outbreak extends beyond the original site/unit

8.14 Enhanced Environmental Cleaning and Disinfection

- See [Section 3.14](#)

Roles and Responsibilities for Confirmed Outbreaks

Only actions and tasks above and beyond those discussed in previous sections are listed here. See [Section 5 Roles and Responsibilities for Confirmed Outbreaks](#) for additional details.

Note: Due to the complex nature of LTC/DSL/Hospice settings, staffing and resident populations, the individual fulfilling the roles and responsibilities within a site may vary from what is outlined below (e.g., some sites may combine the roles of IPC and WHS, or may have designated staff to fulfill these roles)

The Public Health Outbreak Team (MOH, CDC, EPH)

- Provides direction on restrictions to admissions/transfers/discharges to the outbreak unit/facility.
- Provides direction on preventative measures for resident cases.
- Provides direction on restrictions of HCW/staff cases.
- Provides direction on management of unimmunized HCW/staff.
- Provides direction on changes to activities (if applicable) within the unit/facility.
- Advises whether the seasonal influenza vaccine provides adequate protection against the strain of influenza causing the outbreak.
- Directs recommendations for Oseltamivir (Tamiflu) prophylaxis for residents and for unimmunized HCW/staff.

Infection Prevention and Control (Infection Control Professional and/or Infection Control Designate)

Note: In the absence of formal AHS IPC or site ICD coverage, facility administration/manager designates responsibility for these roles. This role may be fulfilled by a single individual at the site, or by a combination of a site Infection Control Designate and supporting groups such as organizational/zone/AHS ICP.

- Directs additional preventative measures required for resident cases.
- Collaborates to implement resident Oseltamivir (Tamiflu) prophylaxis.

- Collaborates to ensure symptomatic residents receive Oseltamivir (Tamiflu) treatment.
- Identifies high risk activities which are recommended to be stopped during the outbreak.

Facility Administration/Facility Management or their Designate

- Supports assessment of resident cases in determining whether site meets outbreak definition.
- Anticipates and manages the impact of HCW/staff exclusion on site operations.
- Ensures that, if recommended, restrictions to admissions / transfer / discharges to the outbreak facility are implemented.
- If facility has restrictions to admissions / transfers / discharges, oversees completion of Risk Assessment Worksheets for operationally necessary resident moves.
- Communicates the status of the facility (open or restricted) to partners and stakeholders.

Site/Unit Manager/Designate

- Provides information about HCW/staff shift patterns (when symptomatic HCW/staff was last onsite) to the Public Health Outbreak Team and OHS / WHS
- Collects information about HCW/staff immunization status. Shares this information with OHS / WHS and the Public Health Outbreak Team.
- Supports assessment of resident cases in determining whether unit meets outbreak definition.
- Anticipates and manages the impact of HCW/staff exclusion on unit operations.
- Ensures that, if recommended, restrictions to admissions / transfer / discharges to the outbreak unit are implemented.
- If unit has restrictions to admissions / transfers / discharges, oversees completion of Risk Assessment Worksheets for operationally necessary resident moves.
- Communicates the status of the unit (open or restricted) to partners and stakeholders.

Occupational Health/Workplace Health and Safety/Designate

***Note:** In the absence of formal AHS WHS or site OHS coverage, facility administration/manager designates responsibility for these roles.*

- Confirms HCW/staff cases and implements HCW/staff exclusions.
- Assesses HCW/staff suitability for return to work.
- Collaborates to determine the immunization status of HCW/staff.
- Identifies HCW/staff who are not fully immunized and communicates work restrictions.
- Implements site plan for how unimmunized HCW/staff are recommended to access and pay for Oseltamivir (Tamiflu) prophylaxis.

Onsite HCW/Staff (hired directly by the site; contracted HCW/staff, AHS HCW/staff)

- Each HCW/staff is responsible for reporting their personal immunization status with Facility Administration/Facility Management and/or Occupational Health/Workplace Health and Safety.
- For unimmunized HCW/staff – Provides decision on whether accepting of Oseltamivir (Tamiflu) prophylaxis or not to Facility Administration/Facility Management and / or OHS / WHS.

Provincial Laboratory for Public Health (ProvLab)

- If applicable, completes additional testing on influenza outbreak specimens.
- Provides guidance regarding match between outbreak strain and strains in the seasonal influenza vaccine.

Attachment 8.1: MOH Notice to all Unimmunized HCW

Exclusion from Work (SAMPLE) Letter used for Confirmed Influenza Outbreaks

The Medical Officer of Health (MOH) or designate has declared an outbreak of influenza at _____ effective _____. Influenza is a serious infectious disease, especially in persons who are elderly or have underlying medical conditions. Workers in health care facilities who come into contact with these vulnerable persons have a duty of care to protect them by being immunized against influenza.

Facility records indicate that you were not immunized against influenza during the _____ influenza season. Under Section 29(2) of the [Public Health Act](#) of Alberta, the MOH has the legal authority to undertake whatever steps are necessary to prevent the spread of a communicable disease to others and may prohibit a person from engaging in their occupation if this activity could transmit an infectious agent.

Because of the risk that you could transmit influenza to vulnerable individuals in your care, effective immediately, the MOH has ordered your manager to **exclude you from further work** in the outbreak facility until:

- a) You receive the influenza immunization now AND commence antiviral prophylaxis for a period of 10 days, or up to a maximum of 14 days dependent on outbreak duration. Protection from immunization takes two weeks to develop completely. Vaccine may be available from pharmacies, Public Health, your family physician, Nurse Practitioner, or your facility. Your family physician, another physician, Nurse Practitioner, or a prescribing pharmacist can also prescribe the appropriate antiviral agent.

OR

- b) You start antiviral prophylaxis immediately WITHOUT receiving influenza immunization. Prophylaxis must be taken for the duration of the outbreak and an initial 10-day supply is recommended to be obtained by prescription from your family physician or through special arrangements at your facility if they exist. Without immunization, you will not develop immunity against influenza, and to continue to work in the event of other influenza outbreaks you will need to take antiviral prophylaxis again.

OR

- c) Two weeks after you have been immunized if you DO NOT take antiviral prophylaxis.

OR

- d) The outbreak is declared over (7 days following onset of symptoms in the last case at the outbreak facility) if you refuse a, b, or c.

You may return to work **immediately** after commencing prophylaxis, provided you do not have any symptoms of influenza (acute onset of headache, chills, and dry cough, followed by fever, muscle aches and pains, runny nose, and/or malaise). If you develop symptoms, the length of time for which you are recommended to stay off work will be recommended by the Public Health Outbreak Team at the time of the outbreak. Generally, a person with influenza is considered infectious for five (5) days.

If you work at health care facilities in addition to the outbreak facility, you may continue to report for work at these facilities if you have complied with either of option (a) or (b) above. If you have not, you are excluded from working in any non-outbreak facility for a period of **three (3) days** after your last shift at the outbreak facility (assuming you remain symptom free), in order to ensure you do not spread influenza to other facilities.

If you have questions about this exclusion, please contact your manager.

Medical Officer of Health

Attachment 8.2 MOH Notice to HCW Immunized less than 14 days prior to declaration of Outbreak

Exclusion from Work (SAMPLE) Letter used for Confirmed Influenza

The Medical Officer of Health (MOH) or designate has declared an outbreak of influenza at _____ effective _____. Influenza is a serious infectious disease, especially in persons who are elderly or have underlying medical conditions. Workers in health care facilities who come into contact with these vulnerable persons have a duty of care to protect them by being immunized against influenza.

Facility records indicate that you were vaccinated against influenza during the _____ Influenza season, but it has been less than 14 days from the date of your immunization and protection from immunization takes two weeks to develop completely. Under Section 29(2) of the *Public Health Act* of Alberta, the MOH has the legal authority to undertake whatever steps are necessary to prevent the spread of a communicable disease to others and may prohibit a person from engaging in their occupation if this activity could transmit an infectious agent.

Because of the risk that you could transmit influenza to vulnerable individuals in your care, effective immediately, the MOH has ordered your manager to **exclude you from further work** in the outbreak facility until:

- a) You commence antiviral prophylaxis. Protection from immunization takes two weeks to develop completely, you therefore must take the prophylaxis until it has been 14 days post-immunization with the current season's influenza vaccine OR for the duration of the outbreak (whichever is shorter). Your family physician, another physician, Nurse Practitioner, or a prescribing pharmacist can prescribe the appropriate antiviral agent.

OR

- b) Two weeks after you have been immunized if you DO NOT take antiviral prophylaxis.

OR

- c) The outbreak is declared over (7 days following onset of symptoms in the last case at the outbreak facility) if you refuse (a).


You may return to work **immediately** after commencing prophylaxis, provided you do not have any symptoms of influenza (acute onset of headache, chills, and dry cough, followed by fever, muscle aches and pains, runny nose, and/or malaise). If you develop symptoms, the length of time for which you are recommended to stay off work will be recommended by the Public Health Outbreak Team at the time of the outbreak. Generally, a person with influenza is considered infectious for five (5) days.

If you work at health care facilities in addition to the outbreak facility, you may continue to report for work at these facilities if you have complied option (a) above. If you have not, you are excluded from working in any non-outbreak facility for a period of **three (3) days** after your last shift at the outbreak facility (assuming you remain symptom free), in order to ensure you do not spread influenza to other facilities.

If you have questions about this exclusion, please contact your manager.

Medical Officer of Health

Attachment 8.3: Influenza Vaccine Decision Tool for HCW/staff



Section 9: Confirmed Gastrointestinal Illness Outbreak

GI Illness Case Definition

At least one of the following criteria must be met and not be attributed to another cause (e.g., *Clostridioides difficile* diarrhea, medication, laxatives, diet, or prior medical condition etc.):

- 2 or more episodes of diarrhea (i.e., loose, or watery stools) in a 24-hour period, above what is normally expected for that individual
- OR**
- 2 or more episodes of vomiting in a 24-hour period
- OR**
- 1 or more episodes of vomiting AND diarrhea in a 24-hour period
- OR**
- 1 episode of bloody diarrhea
- OR**
- Laboratory confirmation of a known enteric pathogen AND at least one symptom compatible with a GI infection e.g., nausea, vomiting, diarrhea, abdominal pain, or tenderness

Note: Laboratory confirmation is not required

GI Illness Outbreak Definition

2 or more cases of GI illness with a common epidemiological link (e.g., same location or same care giver, and evidence of healthcare-associated transmission within the facility), with initial onset within one 48-hour period.

While it is recognized that *Clostridioides difficile* and multi-drug resistant organisms (e.g., MRSA, VRE) can be responsible for clusters or outbreaks, and that some of the measures outlined in this protocol may be applicable in preventing or controlling them, it is beyond the scope of this document to include these organisms, due to their unique epidemiological properties.

See [Section 5](#): General Recommendations for Confirmed Outbreaks.

Outbreaks of infectious GI illness in healthcare facilities can result in high morbidity and a strain on operations. Typically, most of these outbreaks are attributable to norovirus (or viruses that cause similar illnesses, such as sapovirus, rotavirus, astrovirus or adenovirus). Norovirus is extremely communicable, and outbreaks are common. Outbreaks can present in sporadic episodes, or as intensely concentrated events occurring all at once. Attack rates can be quite high (> 50%) in both HCW/staff and residents. Although GI illness outbreaks in healthcare facilities can occur at any time of year, in Alberta most outbreaks occur between October and April.

Most GI illness cases are mild and self-limiting; however, serious dehydration and/or aspiration pneumonia secondary to vomiting can occur in debilitated individuals. Symptoms of GI illness include any combination of nausea, vomiting, diarrhea, and/or abdominal pain, which may be accompanied by myalgia, headache, low-grade fever, and malaise. An outbreak control program is aimed at early detection and elimination of any common sources of exposure. Despite stringent IPC, outbreak control can be difficult. It is vital that infection control measures are implemented promptly, without waiting for laboratory confirmation of an etiologic agent.

Transmission usually occurs via the fecal/oral or vomitus/oral route but can also include contact or droplet spread.

9.1 Infection Prevention and Control Measures

- Consult with ICP/ICD for assistance with IPC issues as required.
- Ensure adequate availability of all supplies as recommended in [Section 3.14](#)
- When a resident GI case is identified, place signage on the outside of the resident's room door indicating that the appropriate precautions are required.
- Appropriate PPE must be worn [Putting on \(Donning\) Personal Protective Equipment](#) and Taking off (Doffing) Personal Protective Equipment (PPE). Instructions are available on the AHS IPC webpage.
- All PPE must be removed, and hand hygiene must be performed before leaving the resident's room.
- Strict hand hygiene is the most important measure in preventing spread of infections for both HCW/staff and residents.
 - Alcohol-based hand rub (ABHR) is the Infection Prevention and Control preferred product for performing hand hygiene, except when providing care for residents/patients with diarrhea and/or vomiting [AHS Hand Hygiene Procedure](#)

Statement on use of Alcohol-based Hand Rub during GI illness Outbreaks

Alcohol-based hand rubs (minimum 60-90% alcohol) are an acceptable alternative to hand washing during GI illness outbreaks when hands are not visibly dirty and when used according to label directions.

Figure 3: Additional Precautions for GI Illness

Implement **Contact Precautions** in addition to Routine Practices for symptomatic residents. Contact Precautions are implemented for symptomatic residents to control the spread of gastrointestinal viruses during GI illness outbreaks (see AHS IPC website Precautions Posters at [Posters | Alberta Health Services](#) for definitions of contact, droplet and droplet and contact precautions).

Implement **Droplet and Contact Precautions if resident is actively vomiting.**

- **Wear clean Gloves** to enter resident room or bedspace when providing direct care to symptomatic residents or when having any contact with items in the resident room, when cleaning an area contaminated with feces or vomitus or gathering/handling specimens.
- **Wear a new Gown** to enter resident room or bedspace when providing direct care to symptomatic residents or when having any contact with items in the resident room' or when cleaning areas contaminated with feces or vomitus to protect against possible contamination of clothing.
- **Wear Eye Protection and a Procedure Mask** to protect your face when there is any risk of sprays of body fluids or when caring for residents who are actively vomiting.

Refer to site IPC guidelines and Doffing poster on the AHS IPC website for instructions [Posters | Alberta Health Services](#).

Maintain at least 2 metres of physical separation between bed/stretchers for contact precautions if diarrhea only.

Maintain at least 2 metres of physical separation between bed/stretchers for droplet and contact precautions if there is active vomiting and diarrhea.

9.2 Administrative Measures

- Advise HCW/staff to report symptoms of GI illness in themselves during the outbreak to the unit/facility Manager, so that their illness can be tracked to capture the scope of the outbreak.
- Case reporting for GI illness will be completed using [Attachment 9.1](#) or zone-specific line lists.

9.3 Resident Restrictions

Resident activities are restricted during GI illness outbreaks; the overarching principle is to prevent congregation of residents in the site during the outbreak. The specific resident restrictions at the outbreak site will be discussed with the Public Health Outbreak Team and partner(s) involved in managing the outbreak. General recommendations are outlined below:

- Whenever possible, symptomatic residents are recommended to be isolated (i.e., remain in their rooms) with meals delivered to them for the duration of the acute illness, and until 48 hours after the last episode of vomiting or diarrhea.
- Symptomatic residents are recommended to only leave the outbreak unit/facility when it is medically necessary; in which case the receiving site is recommended to be alerted that the resident is symptomatic and coming from a facility experiencing a GI illness outbreak and that contact precautions (contact/droplet precautions if vomiting) are to be implemented by the receiving site.

- If an outbreak is confined to a unit, all residents on that unit are recommended to remain on their own unit to avoid contact with other residents in the facility.
- Residents requesting a pass to leave a facility that is under restrictions due to a GI illness outbreak may do so if the resident is asymptomatic. Residents are recommended to be advised that if they become symptomatic while away from their facility, they are recommended to contact their facility, or seek medical attention.
- During an outbreak, consideration is recommended to be given to providing treatment such as physiotherapy or occupational therapy in the resident's room instead of a centralized area; however, residents may be allowed to attend medically necessary activities or appointments provided measures are taken to minimize transmission.

9.4 Restrictions on Affected Units/Site

- Facility/unit status (e.g., open, or restricted admissions) will be determined by the Public Health Outbreak Team at the time the outbreak is declared.
- Decisions regarding GI illness outbreak unit restrictions will be made by the OMT or the Public Health Outbreak Team in consultation with the facility/unit administration.
- Even when admission restrictions are lifted at the recommendation of the OMT or the Public Health Outbreak Team, some residents may still be symptomatic with GI illness. Isolation precautions for symptomatic residents are recommended to remain in effect to prevent further spread of infection.
- The scope of unit restrictions is typically dependent on the extent of the outbreak activity within the facility (one unit, one floor, one wing or the entire facility), the ability to cohort HCW/staff to affected areas and severity of the outbreak (e.g., many residents and HCW/staff affected, new cases continue to develop despite implemented control measures).
- Restrictions typically remain in place until the outbreak has been declared over by the Public Health Outbreak Team or designate. Outbreaks are generally closed following the timeframes below, but may vary due to outbreak assessment on a case-by-case basis, **whichever occurs first:**
 - 48 hours from symptom resolution in the last case
 - OR**
 - 96 hours from onset of symptoms in the last case

Restrictions regarding resident admissions/re-admissions/transfer and activities are **ONLY** modified or lifted by the Public Health Outbreak Team.

If restriction of admissions/transfers is unduly impacting the availability of acute care beds for individuals requiring urgent care, or because of the expressed informed individual resident or family choice (in keeping with AHS's commitment to People Centered Care), the Public Health Outbreak Team will assess the circumstances surrounding the restriction including the degree of risk to the full spectrum of individuals requiring care.

9.5 Admissions/Transfers from Acute Care to an Outbreak LTC, DSL or Hospice Site

- A resident who is hospitalized at another facility prior to the outbreak is not recommended to be transferred back to the facility until the outbreak is declared over. If there are extenuating circumstances, an outbreak facility/site may initiate discussion with the Public Health Outbreak Team to further assess the specific situation.

If an admission/transfer to a site must occur during a GI illness outbreak at the site, following the assessment of the circumstances and consultation with the Public Health Outbreak Team. HCW/staff are recommended to collaborate with the acute care contact before the resident is discharged. The resident is not recommended to be transferred until the site HCW can ensure that the resident/ guardian has information on risks associated with the outbreak and consents to the transfer.

9.6 Transfers from an Outbreak Facility to Acute Care

- If a resident requires acute medical attention or treatment off site (e.g., emergency room, urgent care, dialysis), the outbreak facility must notify the EMS* Dispatcher, the transport staff (EMS crew) and the receiving care facility that the resident is being transferred from a facility experiencing a GI illness outbreak. The transport staff (EMS crew) and the facility receiving the resident can then ensure contact precautions (contact/droplet if resident is vomiting) are in place when the resident arrives there.

*In some communities, [EMS Community Paramedics](#) can provide immediate or scheduled medical supervision and treatments that are currently unavailable in the community setting. A referral is required to access this service.

9.7 Group/Social Activities and other Events

- It is recommended that previously scheduled resident social and special events/activities (e.g., entertainers, school groups, community presentations, and/or communal meals for special holidays) on the affected unit(s) be canceled/postponed for the duration of the outbreak. Consult the Public Health Outbreak Team about group/social activities, or other situations where residents may congregate. If group activities are an essential part of treatment, discuss with the Public Health Outbreak Team.
- Non-resident events (e.g., meetings) previously booked for areas in proximity to areas under restriction in the outbreak facility are recommended to be cancelled or postponed.

9.8 Nourishment Areas / Sharing of Food

- Close the kitchen/nourishment areas accessed by residents/visitors and ensure there is no communal sharing of food in outbreak areas.
- In discussion with the Public Health Outbreak Team, implement measures to minimize resident handling of shared food and surfaces that may touch another resident's food:
 - Close buffet lines, or, have HCW/staff dispense foods from the buffet onto plates for residents.
 - Cease Family-style meal service.
 - Pre-set the tables in common dining areas to minimize resident handling of multiple sets of cutlery.
 - Remove shared food containers from dining areas (e.g., shared pitchers of water, shared coffee cream dispensers, salt & pepper shakers, etc.)
 - For snack programs, dispense snacks directly to residents and use pre-packaged snacks only.
 - If using single service packets of condiments, provide packet directly to each resident, rather than self-serve in a bulk container.
 - Cease activities involving resident participation in food preparation.
 - Other measures as necessary/appropriate

- Ensure that food handling HCW/staff:
 - Practice meticulous hand hygiene
 - Are excluded from work if symptomatic [Section 9.12](#)
- Use of disposable plates and cutlery by symptomatic residents is not required for GI illness outbreak management.
- Normal dishwashing and food preparation area surface sanitizing practices are appropriate during GI outbreaks, i.e., no additional/different disinfection of dishes or surfaces in food preparation areas is needed over and above what is normally done.
- Use dining table coverings that can be easily cleaned and disinfected (i.e., discontinue use of cloth/linen table coverings until the outbreak is over).
- Ensure that all touch surfaces of the tables and chairs (including the underneath edge of the chair seat) are cleaned and disinfected after each use.
- Staff assigned to housekeeping duties are not recommended to be involved in food preparation or food service. Consult the Public Health Outbreak Team with any questions.

9.9 Adult Day Programs in sites Operating Adult Day Programs (e.g., CHOICE/C3)

If the outbreak facility operates an Adult Day Program (e.g., CHOICE/C3), discuss this with the OMT/Public Health Outbreak Team at the time the outbreak is first reported. Generally, the Public Health Outbreak Team will recommend that Adult Day Programs continue to operate in a facility with an ongoing GI illness outbreak **IF**:

- the Adult Day Program is operated in an area physically separate from areas of the facility in which there have been resident cases with GI illness symptoms.
- residents attending the Adult Day Program do not mix with the residents from the outbreak facility.
- Adult Day Program HCW/staff do not provide care in areas of the facility in which there have been outbreak cases.

9.10 Visitor and Designated Family/Support Person(s) (DFSPs)

- Post outbreak signage [Attachment 5.1](#) at the entrance of the facility/unit advising HCW/staff, other professions working in the facility that are not HCW, and visitors of necessary precautions.
- Visitors are recommended to be advised of the potential risk of acquiring illness and advised to practice hand hygiene before and after visiting.
- Advise those who choose to visit during an outbreak to practice good hand hygiene and limit visit to one (1) resident only and exit the facility immediately after the visit.
- Those visiting symptomatic residents must be advised to practice Contact Precautions (contact/droplet precautions if vomiting) to protect themselves.
- Complete restriction of visitation during GI illness outbreaks is typically not recommended by AHS as it may cause emotional hardship to both residents and families. However, if a facility is having difficulty controlling an outbreak, consult the Public Health Outbreak Team.
- Designated Family/Support Person(s) - [See Section 3.10](#)

9.11 Volunteer Restrictions

- Volunteers are recommended to be advised of the potential risk of acquiring illness.
- Advise volunteers of the importance of hand hygiene

- Volunteers who continue to help during an outbreak are managed as outlined in [Section 9.12](#)
- Exclude non-essential volunteers from working in affected areas of the facility (if any can be deemed “non-essential”).

9.12 HCW/Staff Outbreak Measures (Including volunteers, students, physicians)

- HCW/staff are recommended to be advised of the need for daily self-assessment for GI illness symptoms.
- Symptomatic HCW/staff that fit the case definition for GI illness are recommended to contact WHS/OHS and be excluded from work at all care facilities until 48 hours following the last episode of vomiting and/or diarrhea.
- Asymptomatic HCW/staff may work in the outbreak unit as well as other work locations during different shifts.

GI illness among HCW/staff: If there is an unusual increase in GI illness amongst HCW/staff (above the baseline of what would be expected) whether they were present at work with symptoms or not it is recommended to be reported to CEIR as this could be an indicator of a potential outbreak.

9.13 Specimen Collection

- Sites must collect specimens as directed by the Public Health Outbreak Team and arrange for delivery of specimens to the laboratory. It is valuable to collect stool specimens from cases during outbreaks to try and identify the etiology, if possible. Typical GI illness outbreak stool specimen collection procedures are found in [Attachment 3.1](#).

9.14 Enhanced Environmental Cleaning and Disinfection

- Environmental surfaces often become contaminated with feces or vomitus containing viruses or bacteria causing GI illness. Thorough cleaning and disinfection of frequently touched surfaces and equipment can help interrupt disease transmission during GI illness outbreaks. Perform cleaning and disinfection of frequently touched environmental surfaces and equipment in isolation and cohorted areas, as well as high-traffic clinical areas. Frequently touched surfaces include, but are not limited to, commodes, toilets, faucets, hand/bed railing, telephones, door handles, and computer equipment.
- Specific information about environmental cleaning can be found at
 - [Principles for Environmental Cleaning and Disinfection \(albertahealthservices.ca\)](https://albertahealthservices.ca/principles-for-environmental-cleaning-and-disinfection)
 - [Public Health Recommendations for Environmental Cleaning and Disinfection of Public Facilities](#)

Recommended disinfectants

The following disinfectant categories/concentrations are recommended for disinfecting surfaces and equipment during GI illness outbreaks (follow manufacturer’s directions for use):

1. Hypochlorite at a concentration of 1000 parts-per-million. Commercially available hypochlorite-containing solutions are recommended.
2. A surface disinfectant with a Drug Identification Number (DIN) issued by Health Canada with a specific label claim against norovirus, feline calicivirus, or murine norovirus.

An example of a product with this label claim currently in wide use in AHS facilities is 0.5% accelerated hydrogen peroxide. There are other products available with this label claim.

- Equipment is recommended to be cleaned and disinfected only with a product listed in and following the procedures outlined in the manufacturer's directions for that equipment.
- Surfaces must first be cleaned prior to disinfection (2 step process). If the surface disinfectant product used has cleaning properties (detergent/disinfectant) it may be used for both steps, if the surface is visibly clean. Follow manufacturer's directions for use.

Follow recommended cleaning and disinfection protocols (refer to Enhanced Cleaning in [Section 3.14](#) for General Outbreak related Cleaning and disinfection)

- Immediately clean and disinfect areas soiled with emesis or fecal material.
- Use fresh mop head, cloths, cleaning supplies and cleaning solutions to clean affected rooms, and after cleaning large spills of emesis or fecal material.
- Consider discarding all disposable resident-care items and laundering unused linens (e.g., towels, sheets) from resident rooms when the isolation precautions for GI illness are lifted.
- Privacy curtains are recommended to be changed if visibly dirty and when isolation precautions for GI illness are lifted.
- Conduct a thorough, enhanced cleaning in all affected areas at the end of the outbreak.

Note: upholstered furniture and rugs or carpets are recommended to be cleaned and disinfected when contaminated with emesis or stool but may be difficult to clean and disinfect completely. Consult manufacturer's recommendations for cleaning and disinfection of these surfaces. If manufacturer's recommendations are not available, consult the Public Health Outbreak Team. Consider discarding items that cannot be appropriately cleaned/disinfected, when possible/appropriate.

Linen/Laundry

- Appropriate PPE (e.g., gowns) are recommended to be worn if there is a risk of contamination of HCW/staff clothing from body fluids or secretions.
- PPE including gloves are recommended to be removed and hands cleaned once soiled laundry has been placed in the laundry bag.
- If laundry is done in resident laundry rooms (vs. a central laundry room) dedicate one laundry room for soiled laundry from resident's sick with the outbreak illness.
- All linen that is soiled with body fluids are recommended to be handled using the same precautions regardless of the source.
- Remove gross soiling (e.g., feces) with a gloved hand and dispose into toilet. Do not remove excrement by spraying with water.
- Bag or contain soiled laundry at point of care.
- Do not sort or pre rinse soiled laundry in resident care areas.
- Handle soiled laundry with minimum agitation to avoid contamination of surfaces & people. (e.g., roll up)
- Contain wet laundry before placing it in a laundry bag (e.g., wrap in a dry sheet or towel).
- Double bagging is not necessary & not recommended.
- Laundry bags are recommended to be tied securely & not over-filled.

- If the laundry machine has been used to clean laundry soiled with diarrhea or vomiting a bleach cycle of the laundry machine is recommended to be ran (without a load of laundry) before washing other laundry

9.15 Management of “Relapse” Cases

GI illness cases frequently “relapse” (i.e., experience onset of vomiting or diarrhea after being asymptomatic for 24 to 48 hours). The relapse is likely due to malabsorption during an existing norovirus infection rather than being a new infection. These “relapse” GI illness cases:

- are recommended to be isolated until they are free of vomiting and diarrhea for 48 hours, as they may still be infectious.
- are NOT recommended to be counted as new outbreak cases (and therefore NOT be included on daily case listings) – these are not new outbreak cases. Therefore, relapse case(s) alone would not result in the extension of admission restrictions.

Note: If a previously identified GI illness case has onset of GI illness symptoms after being symptom free for **at least seven (7) days**, it is considered a new case.

Attachment 9.1: Data Collection Tool for Gastrointestinal Illness Outbreak Management

It is important that as soon as an outbreak is suspected, front line HCW/staff assess and track symptomatic residents and HCW/staff for surveillance, monitoring, and reporting purposes. Accurately completed lists of cases are recommended to be reported to the Public Health Outbreak Team on a **daily basis** once an outbreak has been declared. The individual responsible for completing and submitting the list of cases are recommended to be determined by site/zone process but may be taken on by site ICP/ICD, unit/facility manager or another responsible HCW/staff in the unit/site. Outbreak data elements that are recommended to be reported daily to the Public Health Outbreak Team include:

Outbreak Facility/Site (name, unit/floor, contact person, phone, and fax)

Date of Report

Population affected at the time outbreak is reported (total resident and HCW/staff population at risk on the outbreak unit/site, number of residents and HCW/staff who meet the case definition)

Outbreak/EI number (as provided by the Public Health Outbreak Team)

Demographics of Cases

- Residents: name, personal health number, date of birth, gender, unit/room number
- HCW/staff: Provide information as requested by PH Outbreak Team: name, gender, occupation, unit they work on

Signs and Symptoms

- Onset date
- Signs and symptoms meeting case definition (vomiting, diarrhea, bloody diarrhea)
- Are clients still on precautions or symptomatic

Lab tests/Results

- Stool specimen (date sent)
- Results

Hospitalization or Death of Cases

- Cases hospitalized (name, personal health number, date of admission, name of hospital)
- Cases who died (name, personal health number, date, and cause of death)

Section 10: Closing an Outbreak

The Public Health Outbreak Team will declare the outbreak over and lift any site restrictions. Restrictions can only be lifted by the Public Health Outbreak Team.

Conduct a thorough, enhanced cleaning in all affected areas at the end of the outbreak as per facility protocols.

Following an outbreak, key program leads need to review and evaluate their role in the outbreak management and revise internal protocols where necessary for improvement. A debriefing may be called by any member of the Outbreak Management Team (OMT) to address outbreak management issues. Depending on type and scale of the outbreak, a summary report including background, details of the investigation, results and recommendations may be written by a member of the OMT and shared with internal/external partners.

If additional residents develop symptoms within 7 days of the outbreak being closed, the Site / Unit is recommended to follow the steps for assessing a potential outbreak in [Section 2](#) and managing a potential outbreak in [Section 3](#) but is recommended to disclose to CEIR that that had recently closed an outbreak.

Appendix A: Sample Letters and Forms for Outbreak Season Preparation

Sample letters

- Letter to Prescribers - [Advance Prescriptions for Antiviral Medication during Influenza Outbreaks \(including Antiviral \(Oseltamivir\) Dosing Recommendations\)](#)
- Letter to Pharmacist - [Provision of Antiviral Medication During Influenza Outbreaks](#)
- [Letter to Designated \(DSL\) and Non-Designated Supportive Living \(NDSL\) Site Operators, AHS Home Care Seniors Health Program Managers/Operation Managers and Case Managers](#)
- [Important Notice to Staff about Influenza and COVID-19](#)

Sample Forms

- [Letter to Resident or Decision Maker](#)
- [Advance Prescription for Oseltamivir \(Tamiflu®\) \(albertahealthservices.ca\)](#)
- [Outbreak Antiviral Prophylaxis in Non-Designated and Designated Supportive Living Site Worksheet \(albertahealthservices.ca\)](#)

Antiviral (Oseltamivir) Dosing Recommendations

Most responsible care providers can access information on influenza antiviral treatment and prophylaxis from the following resources:

- Antiviral use guidelines from Association of Medical Microbiology and Infectious Disease (AMMI) Canada:
[https://www.ammi.ca/Content/Guidelines/Flu%20\(published%20version\)%20FINAL.pdf](https://www.ammi.ca/Content/Guidelines/Flu%20(published%20version)%20FINAL.pdf)
- TAMIFLU® Product Monograph, Roche Canada:
https://www.rochecanada.com/PMs/Tamiflu/Tamiflu_PM_E.pdf
- AHS Healthcare providers can access Lexicomp through Pharmacy Services, Drug Information on AHS Insite

Dosing recommendations for treatment and prophylaxis varies with client age and health (including weight and renal function).

Serum creatinine tests for residents may be required for determining antiviral dosage. Facilities are recommended to prepare for respiratory virus outbreak season each year by ordering serum creatinine and recording resident weights. A baseline temperature is recommended to also be taken and recorded. Ultimately, prescribers are responsible for determining the appropriate antiviral dose for their residents.

Early initiation of antiviral treatment is critical for treatment effectiveness. Providers are recommended to consider whether antiviral treatment can be started using the most recent creatinine clearance estimate for dosing while awaiting blood work and adjusting the timing and dose based on testing results. Most responsible care providers may consider this approach in the following situations:

- renal function has been unstable in the past, or
- patient/resident oral intake/urine output has been poor in the immediate prior period, or
- where creatinine results are older than one year

In the event of antiviral resistance in the outbreak influenza strain, the Zone MOH in combination with the Public Health Outbreak team will make recommendations on the use of antiviral prophylaxis.

Advance Prescriptions for Antiviral Medication during Influenza Outbreaks

Click here to enter a date.

Dear Colleagues,

Provision of antiviral medication during influenza outbreaks for:

- **All residents living in Long Term Care (LTC) and Supportive Living settings**
- **Unimmunized Staff working in LTC and Supportive Living settings**

Influenza viruses circulate throughout our communities every year. In anticipation of influenza outbreaks occurring in LTC and Supportive Living sites, we are requesting your assistance with preparing individuals under your care (i.e., living or working in an LTC or Supportive Living site) for the upcoming influenza season. Supportive Living sites include lodges, manors, seniors' residences, and designated supportive living facilities.

All individuals are recommended to be encouraged to receive their annual influenza immunization. However, even with good immunization rates, outbreaks of influenza commonly occur in sites in which there is communal dining.

OSELTAMIVIR RECOMMENDATIONS DURING INFLUENZA OUTBREAKS

When an influenza outbreak is declared, the Medical Officer of Health (MOH) recommends the following:

- All residents, whether immunized or not, receive oseltamivir antiviral prophylaxis.
- Unimmunized staff (**including unimmunized physicians making site visits**) are required to take oseltamivir antiviral prophylaxis. Staff who are not immunized and are not taking the recommended antiviral prophylaxis are recommended to be excluded from working as outlined in the AHS Guide for Outbreak Prevention and Control in [non-designated supportive living](#) and in [LTC and designated supportive living](#).
- A prescription for a prophylactic dose of oseltamivir for ten (10) days with two (2) refills for five (5) days each is recommended. Antiviral prophylaxis is recommended for 7 days after onset of symptoms of the last resident case. An oseltamivir dosing chart can be found in the Roche Canada Tamiflu product monograph: [\[Product Monograph Template - Standard\] \(rochecanada.com\)](#). An oseltamivir dosing chart, excerpted from the AHS Guide for Outbreak Prevention and Control, is attached as Appendix A of this letter.

ACCESS TO OSELTAMIVIR PROPHYLAXIS - RESIDENTS

If you provide care to residents within an LTC or Supportive Living site:

- Collaborate with the site operator or site administrator to plan for how you will ensure that residents under your care will have timely access to oseltamivir prophylaxis if an influenza outbreak is declared.

If you provide care to Supportive Living residents outside of the site (in the community):

- Residents of Supportive Living sites where the site does not coordinate access to oseltamivir prophylaxis may be asked to contact a prescriber (a physician, nurse practitioner, or prescribing pharmacist) in order to receive an advance prescription for oseltamivir antiviral prophylaxis in preparation for the start of influenza season.
- If a resident requests an oseltamivir prophylaxis advance prescription, please fax a ten (10) day prescription with two (2) refills for five (5) days to the resident's pharmacy. Prescriptions will remain on hand until such time the MOH declares an influenza outbreak at the resident's site or for 1 year (whichever comes first).

ACCESS TO OSELTAMIVIR PROPHYLAXIS – STAFF

- Covenant Health OHS and Alberta Health Services WHS have a process in place for staff requiring antiviral prophylaxis. Staff will be advised to contact their respective OHS/WHS department at the time of an influenza outbreak for assessment and advice.

Non-AHS/Covenant Health staff may be directed to contact a prescriber (a physician, nurse practitioner, or prescribing pharmacist) at the time of an influenza outbreak to obtain a prescription (if indicated).

ALBERTA INFLUENZA ANTIVIRAL DRUG POLICY

The **Alberta Influenza Antiviral Drug Policy as Applied to Vulnerable Populations Living in Congregate Living Settings** defines who is eligible for provincially funded antiviral medication during influenza outbreaks.

- This policy applies to residents of Supportive Living sites
 - Pharmacies that fill prescriptions for antiviral prophylaxis under this policy are compensated under Alberta Blue Cross as per their Pharmacy Benefact; the resident does not have to pay for these medications (antiviral prophylaxis or treatment doses).
- The following individuals are not eligible for publicly funded antiviral medication under this policy:
 - Residents of LTC facilities or nursing homes and patients in a hospital
 - Residents of LTC are covered under the Nursing Homes Act and patients admitted to hospital are covered under the Hospitals Act
 - Health care workers (HCWs) in any setting
 - Although HCWs (employees or volunteers) are not eligible for publicly funded antiviral medication under this policy, unimmunized staff are recommended to still take antiviral prophylaxis. Their employer must have a process in place regarding coverage of cost of antiviral prophylaxis.

If you have questions, please contact (Zone MOH or Zone CDC Nursing) at (contact number).

Thank you for your continued assistance and co-operation.

NAME

Lead Medical Officer of Health - ZONE

Antiviral (Oseltamivir) Dosing Recommendations

Most responsible care providers can access information on influenza antiviral treatment and prophylaxis from the following resources:

- Antiviral use guidelines from Association of Medical Microbiology and Infectious Disease (AMMI) Canada:
[https://www.ammi.ca/Content/Guidelines/Flu%20\(published%20version\)%20FINAL.pdf](https://www.ammi.ca/Content/Guidelines/Flu%20(published%20version)%20FINAL.pdf)
- TAMIFLU® Product Monograph, Roche Canada:
https://www.rochecanada.com/PMs/Tamiflu/Tamiflu_PM_E.pdf
- AHS Healthcare providers can access Lexicomp through Pharmacy Services, Drug Information on AHS Insite

Dosing recommendations for treatment and prophylaxis varies with client age and health (including weight and renal function).

Serum creatinine tests for residents may be required for determining antiviral dosage. Facilities are recommended to prepare for respiratory virus outbreak season each year by ordering serum creatinine and recording resident weights. A baseline temperature is recommended to also be taken and recorded. Ultimately, prescribers are responsible for determining the appropriate antiviral dose for their residents.

Early initiation of antiviral treatment is critical for treatment effectiveness. Providers are recommended to consider whether antiviral treatment can be started using the most recent creatinine clearance estimate for dosing while awaiting blood work and adjusting the timing and dose based on testing results. Most responsible care providers may consider this approach in the following situations:

- renal function has been unstable in the past, or
- patient/resident oral intake/urine output has been poor in the immediate prior period, or
- where creatinine results are older than one year

In the event of antiviral resistance in the outbreak influenza strain, the Zone MOH in combination with the Public Health Outbreak team will make recommendations on the use of antiviral prophylaxis.

Provision of Antiviral Medication During Influenza Outbreaks

[Click here to enter a date.](#)

Dear Community Pharmacist,

Re: Provision of antiviral medication during influenza outbreaks for:

- **All residents living in Long Term Care (LTC) and Supportive Living settings including lodges, manors, seniors' residences, and designated supportive living facilities**
- **Unimmunized staff working in LTC and Supportive Living settings**

Over the next several weeks, in preparation for the upcoming influenza season, your pharmacy may be receiving advance influenza antiviral prophylaxis prescriptions from prescribers for residents living in LTC and Supportive Living sites. **Please do not dispense these advance prescriptions until you are notified of a laboratory-confirmed influenza outbreak by the Alberta Health Services (AHS) (insert zone) Zone Medical Officer of Health (MOH) or Communicable Disease Control (CDC) nursing team, or by the facility.**

When an influenza outbreak is declared at a congregate living site, the MOH recommends that all residents, whether immunized or not, receive oseltamivir antiviral prophylaxis. Unimmunized staff are also required to take antiviral prophylaxis if they are working during the outbreak. You may receive notification of a confirmed outbreak directly from a facility or from the (insert zone) MOH or CDC Outbreak Team. **In the event you do not receive this notification, you must contact the outbreak facility directly and speak to the outbreak site designate to gather the necessary information.**

- Following a notification of a laboratory-confirmed influenza outbreak at a site, please dispense advance prescriptions for antiviral prophylaxis as soon as possible, and deliver resident antiviral medications, as appropriate.
- At the time of an influenza outbreak, you may receive prescriptions for antiviral **prophylaxis** for residents **without** advance prescriptions and for unimmunized staff working at the outbreak site. You may also receive prescriptions for antiviral **treatment** for symptomatic residents.

Alberta Influenza Antiviral Drug Policy

The **Alberta Influenza Antiviral Drug Policy as Applied to Vulnerable Populations Living in Congregate Living Settings (CLS)** defines who is eligible for provincially funded antiviral medication during influenza outbreaks.

- **This policy applies to residents of Supportive Living sites only**
 - Pharmacies that fill prescriptions for antiviral prophylaxis under this policy are compensated under Alberta Blue Cross as per their Pharmacy Benefact; the resident does not have to pay for antiviral medication (prophylaxis or treatment doses).
- **The following individuals are not eligible for publicly funded antiviral medication under this policy:**
 - Residents of LTC facilities (including nursing homes and auxiliary hospitals) and patients in a hospital
 - Residents of LTC are covered under the Nursing Homes Act and patients admitted to hospital are covered under the Hospitals Act
 - Health care workers (HCWs) in any setting
 - Although HCWs (employees or volunteers) are not eligible for **publicly funded** antiviral medication, unimmunized staff are recommended to still take antiviral

prophylaxis if they are working during the outbreak. Their employer will advise their employees regarding coverage of the cost of antiviral prophylaxis.

Prescribing Pharmacists

- Supportive Living sites may look to collaborate with a prescribing pharmacist to obtain antiviral prophylaxis prescriptions.
- If you are a prescribing pharmacist, you may receive requests from Supportive Living sites or from individual residents to assist with influenza antiviral prophylaxis advance prescriptions or with antiviral prophylaxis prescriptions at the time an influenza outbreak is declared.

Important Things to Note

- Any resident without an Alberta Personal Health Number is covered by this policy (**Please refer to the Pharmacy Benefact for additional information**). Please ensure you are a party to the Alberta Blue Cross Pharmacy Agreement prior to submitting the prescribed Antivirals Dispensing claims.
- Refer to the current Tamiflu (Roche Canada) product monograph for Tamiflu dosing recommendations.
- Covenant Health OHS and Alberta Health Services WHS have a process in place for staff requiring antiviral prophylaxis. Covenant and AHS staff will be advised to contact their respective OHS/WHS department at the time of an influenza outbreak for assessment and advice. Staff working at all other LTC, and Supportive Living facilities are recommended to contact their OHS/WHS designate.
- **If you have questions about billing, you must contact Alberta Blue Cross directly.**

If you have clinical questions, please contact (Zone MOH or Communicable Disease Control) at (contact number for MOH or CDC). When calling, please indicate that you are calling about **(insert zone)** Zone advance antiviral prophylaxis prescriptions for influenza outbreaks.

Thank you for your continued assistance and co-operation.

ADD MOH Signature

Name, designation
Lead Medical Officer of Health – Zone
Alberta Health Services

Outbreak Preparation Letter

To: All DSL, NDSL Site Operators, AHS Home Care Seniors Health Program
Managers/Operation Managers & Case Managers

Date:

Subject: Outbreak Preparation for 2022/23 Season

Dear Operators and Seniors Health Staff:

Outbreaks of respiratory illness (including influenza and [COVID-19](#)) and gastrointestinal illness are possible in Supportive Living facilities. Preparing for and responding to outbreaks is a shared responsibility.

Alberta Health Services Public Health kindly requests your assistance in preparing for the 2022-2023 outbreak season. These tasks will help to protect both staff and residents and will facilitate an effective response if an outbreak occurs at your site.

1. Ensure that all residents have a prescription for influenza antiviral prophylaxis (Oseltamivir [Tamiflu]) available in-advance of, or within a very short time of, an outbreak being confirmed.
 - a. **Arrange for advance prescriptions:**
 - i. **DSL Site Operators in collaboration with AHS Home Care:** Options to ensure advance prescriptions are in place include:
 1. Arranging for resident influenza prophylaxis prescriptions through each resident's most responsible health practitioner (a physician, prescribing pharmacist, or nurse-practitioner). OR
 2. Working with or contracting a single prescriber for the whole site (a physician, prescribing pharmacist, or nurse-practitioner).
 - ii. **NDSL Site Operators in collaboration with Residents:** Options to ensure advance prescriptions are in place include:
 1. Working with or contracting a single prescriber for the whole site (a physician, prescribing pharmacist, or nurse-practitioner). OR
 2. Advising residents to see their primary care provider or another community prescriber (a physician, prescribing pharmacist, or nurse practitioner) to request an advance prescription. Please distribute to residents or guardians/substitute decision-makers the *Resident Outbreak Preparation Letter* and the [Advance Prescription for Oseltamivir \(Tamiflu\)](#) prescription template, within the next month. Please continue to distribute the letter for future admissions until April 30, 2023.
 - b. **Make a plan for how to respond at the time of an outbreak:** Contact a local prescriber (a physician, prescribing pharmacist, or nurse-practitioner) to make arrangements in preparation for immediate access to antivirals for all residents who did not obtain an advance prescription at the time of an outbreak notification. For your convenience, a prescribing pharmacist can be found at this link: [Alberta College of Pharmacy \(abpharmacy.ca\)](https://abpharmacy.ca)

2. Consider creating and maintaining an up-to-date list of residents throughout the outbreak season until April 30, 2023, using the [Outbreak antiviral prophylaxis in non-designated and designated supportive living sites worksheet](#). We suggest the worksheet be completed in advance of an outbreak to ensure timely and effective management at the facility. Please note the following:
 - a. Information about whether residents have received annual influenza vaccine and the number of COVID-19 vaccine doses received may be requested by CDC nursing.
 - b. When an influenza outbreak occurs, residents who do not have an advance prescription for antiviral prophylaxis will need to see a prescriber to get a prescription as soon as possible.
 - c. Information about whether residents have received influenza antiviral prophylaxis may be requested by CDC nursing at the time of an influenza outbreak.
3. Recommend that staff receive all doses of [COVID](#)-19 vaccine that they are eligible for (including boosters) and an annual dose of seasonal influenza vaccine. Provide a copy of the letter *Important Notice to Staff About Influenza and COVID-19* to all staff, including new hires, until April 30, 2023.
4. Recommend that residents receive all doses of [COVID](#)-19 vaccine that they are eligible for (including boosters) and an annual dose of seasonal influenza vaccine.
5. Develop a site plan for how to access specimen collection supplies at the time of an outbreak.

Although roles and responsibilities can vary between sites, site operators, AHS Home Care Managers and Case Managers are recommended to collaborate to ensure that each site is prepared for the upcoming outbreak season.

Outbreak Management Resources:

Outbreak management recommendations and resources changed frequently in the past outbreak season and will likely continue to change frequently during the upcoming season. For the most up to date information, please review the following resources frequently:

- Zone MOH website: (zone specific link)
- AHS Infection Prevention & Control (IPC) Outbreak Management: <https://www.albertahealthservices.ca/ipc/Page6421.aspx>
- Guide for Outbreak Prevention and Control in Non-Designated Supportive Living Sites: <https://www.albertahealthservices.ca/assets/healthinfo/flu/hi-flu-care-and-treat-guidelines.pdf>
- Guide for Outbreak Prevention and Control in Long Term Care, Designated Supportive Living and Hospice Sites: <https://www.albertahealthservices.ca/assets/healthinfo/flu/hi-flu-prov-hlsl.pdf>
- [AHS Health Professionals COVID-19 information:](#) <https://www.albertahealthservices.ca/topics/Page16947.aspx>

Questions regarding outbreak preparation and response can be directed to (zone specific contact information).

Thank you very much for your partnership in outbreak management!

Name, designation
Lead Medical Officer of Health – Zone
Alberta Health Services

SAMPLE

Important Notice to Staff about Influenza and COVID-19

Date

Why is it important to receive a complete COVID-19 immunization series (including any recommended boosters)?

We all must do our part to protect one another. Immunization is the single most effective means of protecting yourself, your loved ones, and the greater community from COVID-19. Without immunization, Albertans are at risk of developing severe illness and even death from this virus. Vaccines strengthen your immune system by building antibodies to help prevent diseases. Immunization is safe. It is still recommended to get the vaccine even if you were previously infected.

Why is it important for you to be immunized with influenza vaccine every year?

Getting immunized against influenza every year is the most effective way to prevent the spread of the virus. Healthcare workers (HCWs) have a unique responsibility to protect their own health as well as the wellbeing of those around them who may be at risk. Other reasons to get immunized include: 1) HCWs are at a higher risk of exposure to influenza than adults in non-healthcare work settings; 2) HCWs may transmit influenza to vulnerable patients, and their own family and friends; and 3) Immunized HCWs decreased risk of illness, serious outcomes and even death for themselves and others.

If you do not get immunized and an influenza outbreak occurs, what happens?

In the event of an influenza outbreak at your site, the <insert zone>Medical Officer of Health may make recommendations that will impact unimmunized staff members. **This may include being excluded from work or starting a prescription antiviral medication called Oseltamivir for the duration of the outbreak.** Depending on employer/employee health insurance arrangements, you may be required to pay for the antiviral.

For more information, please speak with your manager or call Health Link Alberta at 811.

Visit AHS websites: COVID-19 information - <https://www.albertahealthservices.ca/covid19>
Influenza information - <https://www.albertahealthservices.ca/influenza>

Thank you for your attention to this important matter.

Sincerely,

Medical Officers of Health
Alberta Health Services, Zone

Resident or Alternate Decision Maker Fall Letter

Dear Resident or Alternate Decision Maker,
Date

Fall is the time to prepare for respiratory outbreak season. An outbreak is an increase in the number of people who live at a facility and become sick with the same kind of infection. Individuals living in lodges and supportive living facilities may be exposed to different kinds of respiratory infections including influenza and COVID-19. Those who are age 65 or older and anyone with chronic medical conditions are most at risk of serious illness and hospitalization.

Outbreaks may occur at the facility you live in. The table below shows some ways to keep yourself from becoming sick, and to protect yourself if there is an outbreak.

How to protect yourself from infections
<ul style="list-style-type: none"> • Wash your hands often and thoroughly with soap and water or use an alcohol-based hand rub. • Practice physical distancing (2 meters/6 feet) when in public places. • Stay in your room when you are sick. Tell someone in your facility when you are feeling sick. • Consider wearing a mask in public places. • Cough or sneeze into your sleeve or a tissue - not your hands. • Consider getting immunized to protect yourself against vaccine preventable illness [e.g., Pneumococcal]
How to protect yourself from influenza
<ul style="list-style-type: none"> • Get your annual influenza vaccine every fall. • A prescriber (physician, nurse practitioner, or prescribing pharmacist) can give you a prescription for a medicine called Oseltamivir (Tamiflu®) that can help to keep you from getting ill if there is an influenza outbreak at your site. See a prescriber as soon as possible to get a prescription in advance. Bring the attached form called <i>Advance Prescription for Oseltamivir (Tamiflu®)</i> with you. <ul style="list-style-type: none"> ○ This medication is filled and taken during an outbreak even if you have been immunized against influenza as an extra level of protection. ○ There is no charge for this medication.
How to protect yourself from COVID-19
<ul style="list-style-type: none"> • Get all recommended doses of COVID-19 vaccine. • For the most updated information please visit: https://www.albertahealthservices.ca/topics/Page16944.aspx

If you have further questions, please contact the manager at your site.

Thank you for your attention to this important health matter.

Medical Officer of Health

Appendix B: Symptoms? What Do I Do to Prevent a Potential Outbreak

This resource was developed to help facilities/sites manage symptomatic residents before the Reporting Definition is met (prior to calling CEIR).

- This resource can be printed and made available to the front-line staff for daily use. Print copies directly from the web page online - [Outbreak Management | Alberta Health Services](#)


Symptoms? What Do I Do to Prevent a Potential Outbreak?

Prevent a Potential Outbreak in Long-Term Care, Designated Supportive Living, Hospice Sites

All staff share the responsibility for keeping residents safe. Start the following early actions when you identify a resident with new symptoms. Implementing these early actions can prevent a potential outbreak. Refer to the section identified below in the [Guide for Outbreak Prevention and Control in LTC/DSL/Hospice Sites](#) for further information.

	Guide Section
1. Manage the symptomatic resident (if applicable)	
<input type="checkbox"/> Isolate the symptomatic resident in their room on appropriate precautions .	3.1/3.3
<input type="checkbox"/> Assess whether the symptomatic resident meets the Respiratory Surveillance Case Definition in Table A.	2.1
<input type="checkbox"/> Assess whether the symptomatic resident meets the Gastrointestinal (GI) Illness Outbreak Reporting Definition in Table B.	2.1
<input type="checkbox"/> Follow site process to arrange for testing for symptomatic resident if required for medical management.	2.1
<input type="checkbox"/> Post signs (precautions, donning/doffing) outside of the room of symptomatic residents.	3.1
<input type="checkbox"/> Arrange for symptomatic resident to receive meal service in their room.	3.8
<input type="checkbox"/> Gather appropriate supplies - hand hygiene products, PPE, linen, and specimen collection supplies.	3.2
<input type="checkbox"/> All staff are to wear appropriate PPE - see Infection Prevention and Control Risk Assessment (IPCRA)	3.1
<input type="checkbox"/> Use strict hand hygiene at all times.	3.1
<input type="checkbox"/> Follow all IPC measures.	3.1/3.14
2. Communicate	
<input type="checkbox"/> Contact manager/supervisor to report symptomatic or positive residents.	2.2

Date: March 2023 Version 2
Created by: [Notifiable Disease & Outbreak Management | Alberta Health Services](#)
Page: 1/2



Appendix C: Outbreak Checklists

These new resources were developed to help facilities manage outbreaks by outlining the key tasks or responsibilities that are recommended to be considered when an outbreak is opened. The checklists are to be used in combination with the guide so the relevant guide section is indicated on the right-hand side indicating where more information and detail can be found. These checklists are recommended to be used in combination with site-specific protocols as applicable.

- The four different checklists are:
 - COVID-19
 - Influenza-like Illness
 - Influenza
 - Gastrointestinal
- The checklists can be printed and made available to the front-line staff for use.
- Below is an example of the COVID-19 checklist. All the checklists can be printed directly from the webpage online - [Outbreak Management | Alberta Health Services](#)

COVID-19 Outbreak Management

Manage a confirmed outbreak in Long-Term Care, Designated Supportive Living, Hospice Sites

All staff share the responsibility for keeping residents safe. The following actions should be completed once a COVID-19 outbreak has been confirmed. This checklist provides general measures only – always refer to the corresponding section in the [Guide for Outbreak Prevention and Control in Long Term Care, Designated Supportive Living, and Hospice Sites](#) for detailed recommendations. When applicable, follow site-specific protocols.

1. Infection Prevention Control Measures	Guide Section
<input type="checkbox"/> Monitor to identify newly symptomatic residents at the site – see Table A.	2.1/3.2
<input type="checkbox"/> Isolate residents with symptoms.	3.1
<input type="checkbox"/> Place signage outside the room of symptomatic residents, alerting HCW/staff and visitors of precautions required.	3.1/5.2
<input type="checkbox"/> Conduct an Infection Prevention and Control Risk Assessment (IPCRA) prior to every resident interaction.	6.1
<input type="checkbox"/> Wear continuous mask and eye protection in all resident care areas. Additional PPE may be required based on an Infection Prevention and Control Risk Assessment (IPCRA) .	5.2
<input type="checkbox"/> Perform strict hand hygiene in accordance with AHS Hand Hygiene Policy or site policy.	3.1
2. Administrative Measures	Guide Section
<input type="checkbox"/> Notify appropriate HCW/staff/departments within the site facility.	5.2
<input type="checkbox"/> Advise HCW/staff of relevant work recommendations and/or restrictions.	5.2
<input type="checkbox"/> Place outbreak signage at the entrance of the facility/unit.	5.2
<input type="checkbox"/> Inform residents' families/guardians/agents of the facility outbreak status.	5.2
<input type="checkbox"/> Report daily using Facility CDC Outbreak Daily Report Portal (and to IPC as per Zone process)	6.2

Date: March 2023 Version 2
 Created by: [Notifiable Disease & Outbreak Management | Alberta Health Services](#)
 Page: 1/4

