COVID-19 daily epidemiology update

Updated: August 10, 2021, 7 pm EST


Summary of COVID-19 cases across Canada and over time. Contains detailed data about the spread of the virus over time and in different regions of the country. Includes breakdowns by age and sex or gender. Provides an overview of hospitalizations and deaths, testing, variants of concern and exposures.

Key updates as of August 10, 2021, 7 pm EST

Cases today Total cases Active cases Total resolved Deaths today Total deaths 1,346 1,443,433 12,548 1,404,202 6 26,683 Daily percent positive (last 7 days) Daily tests per 100,000 population (last 7 days) Total tests performed 39,070,051 2.4% 151

- We update these sections daily at 7:00 PM EST: Key updates, Current situation and National overview.
 Laboratory data represents specimens received by labs up to August 8, 2021 to allow time to process results.
- We update these sections every Friday: Epidemic curve, Demographics, How people were exposed, and Severe illness and outcomes.
- Most cases (64.7%) and deaths (77.4%) were reported by Ontario and Quebec.
- Of the 11 jurisdictions reporting updates, no new cases were reported in no provinces and territories in the past 24 hours.
- Of the 11 jurisdictions reporting updates, no new deaths were reported in 8 provinces and territories in the past 24 hours.

Current situation



The count of cases (last 7 days) of COVID-19 in Canada was 7,795 as of August 10, 2021.

This information is based on data our provincial and territorial partners published on cases, deaths, and testing daily, and are current as of the day they are published. Today's numbers are current as of August 10, 2021, 2021, 7 PM EST. For the most up to date data for any province, territory or city, please visit their website. The number of cases or deaths reported on previous days may differ slightly from those on the provincial and territorial websites as these websites may update historic case and death counts as new information becomes available.

Areas in Canada with cases of COVID-19 as of August 10, 2021

| | Total cases | 3 | Cases I | ast 7 | Active c | ases | Resolved | Deaths | | Deaths days | last 7 | Total tests performed | Moving a tests per last 7 da | formed | Moving average positivity last 7 days |
|------------------------------|-------------|-------|---------|-------|----------|-------|-----------|--------|-------|----------------|--------|-----------------------|------------------------------------|--------|--|
| Location | Count | Rate* | Count | Rate* | Count | Rate* | Count | Count | Rate* | Count | Rate* | Count | Count | Rate* | Percent |
| Canada | 1,443,433 | 3,798 | 7,795 | 21 | 12,548 | 33 | 1,404,202 | 26,683 | 70 | 67 | 0 | 39,070,051 | 57,235 | 151 | 2.4% |
| British Columbia | 153,313 | 2,978 | 2,185 | 42 | 3,321 | 65 | 148,215 | 1,777 | 35 | 1 | 0 | 3,176,702 | 9,582 | 186 | 4.2% |
| Alberta | 237,306 | 5,367 | 1,754 | 40 | 3,463 | 78 | 231,513 | 2,330 | 53 | 3 | 0 | 4,915,363 | 6,377 | 144 | 4.8% |
| Saskatchewan | 50,679 | 4,300 | 379 | 32 | 634 | 54 | 49,461 | 584 | 50 | 6 | 1 | 971,341 | 1,548 | 131 | 5.3% |
| Manitoba | 57,891 | 4,197 | 175 | 13 | 580 | 42 | 56,127 | 1,184 | 86 | 6 | 0 | 933,742 | 1,627 | 118 | 2.1% |
| Ontario | 553,125 | 3,754 | 1,563 | 11 | 2,494 | 17 | 541,222 | 9,409 | 64 | 49 | 0 | 16,590,360 | 18,171 | 123 | 1.6% |
| Quebec | 379,673 | 4,428 | 1,617 | 19 | 1,909 | 22 | 366,523 | 11,241 | 131 | 1 | 0 | 10,479,329 | 16,060 | 187 | 1.5% |
| Newfoundland and Labrador | 1,443 | 276 | 6 | 1 | 13 | 2 | 1,423 | 7 | 1 | 0 | 0 | 315,588 | 210 | 40 | 0.4% |
| New Brunswick | 2,447 | 313 | 57 | 7 | 65 | 8 | 2,336 | 46 | 6 | 0 | 0 | 399,332 | 746 | 95 | 0.9% |
| Nova Scotia | 5,908 | 603 | 14 | 1 | 17 | 2 | 5,798 | 93 | 10 | 0 | 0 | 1,053,453 | 2,701 | 276 | 0.1% |
| Prince Edward Island | 214 | 134 | 6 | 4 | 6 | 4 | 208 | 0 | 0 | 0 | 0 | 180,954 | 178 | 111 | 0.4% |
| Yukon | 633 | 1,505 | 36 | 86 | 44 | 105 | 581 | 8 | 19 | 1 | 2 | 9,129 | N/A | N/A | N/A |
| Northwest Territories | 131 | 290 | 3 | 7 | 2 | 4 | 129 | 0 | 0 | 0 | 0 | 26,020 | 26 | 57 | 0.6% |
| Nunavut | 657 | 1,670 | 0 | 0 | 0 | 0 | 653 | 4 | 10 | 0 | 0 | 18,662 | 9 | 24 | 0.0% |

^{*} Rate per 100,000 population


Figure 1b. Moving average of


cases (last 7 days)

of COVID-19 in Canada as of

August 10, 2021, 7 pm EST

The figures below show cases over time. The range of dates (January 31st, 2020 - present date) is the same for each figure. This allows you to compare the provinces and territories on the same timescale.





This information is based on data our provincial and territorial partners published on cases, deaths, and testing daily, and are current as of the day they are published. Today's numbers are current as of August 10, 2021, 7 PM EST. For the most up to date data for any province, territory or city, please visit their website. The number of cases or deaths reported on previous days may differ slightly from those on the provincial and territorial websites as these websites may update historic case and death counts as new information becomes available.

Downloadable data (in .csv format).

Note: Out of the total number of people tested, 76 were repatriated travellers, of which 13 were cases.

National overview

There have been over **39,070,051** COVID-19 tests performed in Canada or **457,522 tests per 1 million people**. For information about testing trends, please see the <u>Detailed weekly epidemiological report (PDF)</u>.

Table 1. Daily* change in the number of cases, deaths and tests performed, by province or territory, as of August 10, 2021, 7 pm EST

| Location | New cases | New deaths | Tests performed |
|---------------------------|-----------|------------|-----------------|
| Canada | 1,346 | 6 | 45,617 |
| British Columbia | 395 | 0 | 6,947 |
| Alberta | 279 | 2 | 4,464 |
| Saskatchewan | 65 | 2 | 1,301 |
| Manitoba | 31 | 0 | 1,271 |
| Ontario | 321 | 2 | 15,805 |
| Quebec | 234 | 0 | 13,078 |
| Newfoundland and Labrador | N/A | N/A | 161 |
| New Brunswick | 8 | 0 | 753 |
| Nova Scotia | 1 | 0 | 1,713 |
| Prince Edward Island | 2 | 0 | 113 |
| Yukon | 9 | 0 | N/A |
| Northwest Territories | 1 | 0 | 6 |
| Nunavut | N/A | N/A | 5 |

^{*} The new cases, deaths and tests reflect the difference between a province or territory's current report and their last report. Some provinces and territories do not update daily.

N/A means that no daily update was provided by the province or territory.

COVID-19 variants in Canada

All viruses, including COVID-19, change over time. These changes are called mutations, and result in variants of the virus. Not all mutations are of concern. Most do not cause more severe illness. However, some mutations result in variants of concern or variants of interest.

A variant of concern has mutations that are significant to public health. Before a variant of interest is considered one of concern, scientists and public health professionals must determine if the mutations result in an actual change in the behaviour of the virus. For example, it might:

- · spread more easily
- cause more severe illness
- · require different treatments, or
- · reduce vaccine effectiveness

There are several variants of interest that have mutations similar to variants of concern, but we don't yet know if they pose a higher risk to public health.

The Public Health Agency of Canada (PHAC) works with provincial and territorial partners and the Canadian COVID-19 Genomics Network (CanCOGeN) to sequence a percentage of all positive COVID-19 test results. Sequencing reveals the genetic code of the virus, which tells us which variant is involved in a specific case of COVID-19. We report the proportion of COVID-19 variants in Canada every week.

We collect evidence to determine if new variants meet the definition for a <u>variant of concern or a variant of interest</u>. Many variants are being tracked across Canada and around the world. Variants of concern now represent a majority of COVID-19 cases in Canada.


Four VOC (Variants of concern)s have been detected in most provinces and territories:

- B.1.1.7 (Alpha)
- B.1.351 (Beta)
- P.1 (Gamma)
- B.1.617.2 (Delta)

New variants will continue to appear. We must remain vigilant and take all available measures to limit spread.

Figure 2. Weekly variant breakdown Updated: August 6, 2021, 4 pm EST

The graphic shows the percentage mix of COVID-19 variants detected in Canada through whole genome sequencing, by week of sample collection. You can see the numbers for each date by hovering over, tabbing to, or long-pressing any of the bars. To see a specific variant or variant grouping, click or press return. Repeat to restore the complete graph.



Week of sample collection

This information is based on whole genome sequencing from surveillance testing in all provinces and territories. In addition to sequencing done by the National Microbiology Laboratory in Winnipeg, data is included from <u>provincial and territorial</u> laboratories.

Sequencing takes from 1 to 3 weeks to complete, so the proportions for recent weeks may change as more data are added. Surveillance in each province or territory is organized and prioritized according to local needs and may change from time to time. Because of differences in local sampling and reporting, the percentages illustrate trends rather than precise measurements.

Weekly variant breakdown

Percentage of COVID-19 cases identified through whole genome sequencing, presented by variant and by week of sample collection.

| Variant Grouping | May 09, 2021 (n=4,008) | May 16, 2021 (n=3,477) | May 23, 2021 (n=2,799) | May 30, 2021 (n=2,897) | Jun 06, 2021 (n=2,926) | Jun 13, 2021 (n=2,962) | Jun 20, 2021 (n=2,401) | Jun 27, 2021 (n=1,100) | Jul 04, 2021 (n=771) | Jul 11, 2021 (n=105 |
|----------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|--|--|----------------------------|----------------------------|
| Variants of concern | 89.8% | 93.2% | 92.5% | 94.6% | 90.2% | 89.9% | 90.6% | 95.4% | 94.5% | 83.0% |
| Alpha | 55.5% | 53.1% | 53.7% | 55.7% | 51.6% | 41.2% | 45.1% | 18.9% | 28.1% | 1.0% |
| Beta | 1.0% | 0.9% | 0.3% | 0.8% | 0.2% | 0.3% | 0.9% | - | 0.4% | 1.0% |
| Delta | 8.3% | 9.6% | 13.6% | 16.3% | 22.7% | 35.3% | 33.9% | 68.7% | 62.1% | 78.1% |
| Gamma | 25.0% | 29.6% | 24.9% | 21.8% | 15.7% | 13.1% | 10.7% | 7.8% | 3.9% | 2.9% |
| Variants of interest | 3.9% | 3.0% | 3.5% | 2.9% | 3.8% | 3.5% | 3.1% | 1.9% | 0.2% | 5.7% |
| A.23.1 | 0.1% | - | - | - | - | - | - | - | - | - |
| B.1.1.318 | 0.4% | 0.7% | 0.5% | 0.9% | 1.3% | 0.8% | 1.0% | 0.9% | 0.1% | - |
| B.1.617.3 | - | - | - | - | - | 0.1% | - | - | - | _ |
| Epsilon | 0.7% | 0.9% | 0.4% | 0.5% | 0.1% | 0.1% | 0.1% | - | - | _ |
| Eta | 1.6% | 1.1% | 1.1% | 0.4% | 1.9% | 2.1% | 1.8% | 0.2% | 0.1% | - |
| lota | 0.8% | 0.1% | 1.5% | 1.0% | 0.5% | 0.4% | 0.2% | 0.8% | - | - |
| Карра | 0.2% | 0.2% | - | 0.1% | _ | - | - | _ | - | - |
| Lambda | 0.1% | _ | - | - | _ | - | - | _ | - | 5.7% |
| Theta | - | - | - | - | _ | - | - | _ | - | - |
| Zeta | - | _ | - | - | _ | - | - | _ | - | - |
| Other variants | 5.2% | 3.6% | 3.9% | 2.4% | 5.6% | 6.3% | 5.7% | 2.8% | 5.1% | 11.6% |
| B.1.438 | 3.5% | 1.9% | 1.4% | 0.4% | 3.7% | 3.8% | 2.0% | 0.5% | 0.6% | - |
| B.1.621 | - | 0.1% | 0.3% | 0.1% | 0.5% | 0.4% | 0.6% | 0.9% | _ | 3.8% |
| Other | 1.7% | 1.6% | 2.2% | 1.9% | 1.4% | 2.1% | 3.1% | 1.4% | 4.5% | 7.8% |
| | | | | | | | | | | |

Contributing laboratories:

- Public Health Ontario (PHO)
- · Newfoundland and Labrador Eastern Health
- New Brunswick Vitalité Health Network
- McMaster University
- Manitoba Cadham Provincial Laboratory
- Laboratoire de santé publique du Québec (LSPQ)
- BCCDC Public Health Laboratory
- Alberta Precision Labs (APL)
- National Microbiology Laboratory (NML) supplemental sequencing for all provinces and territories

Detailed case information

The tables and figures below reflect detailed case information provided to the Public Health Agency of Canada (PHAC) by health authorities in the provinces and territories. This data is updated every week. It may change as we get more information about cases.

Updated: August 6, 2021, 7 pm EST

Epidemic curve

As of August 6, 2021, 7 pm EST, PHAC has received detailed case report data on 1,435,035 cases. Both exposure and symptom onset date were available for 1,070,583 (74.6%) cases $\frac{1}{2}$.

The shaded area on the far right of Figure 3 represents a period of accumulating data. This is the period of time (1 to 2 weeks) before the latest cases are reported to PHAC. This delay is a result of the time required to seek health care, get tested and receive results. It also takes time for public health authorities to gather information on cases. We update this information as it becomes available.

Figure 3. COVID-19 cases (n=1,435,035 1) in Canada by date of illness onset 2 as of August 6, 2021, 7 pm EST (total cases)




Figure 3. COVID-19 cases (n=1,070,583 1) in Canada by date of illness onset 2 as of August 6, 2021, 7 pm EST (by exposure)






Figure 3. COVID-19 cases (n=1,434,642 $\frac{1}{2}$) in Canada by date of illness onset $\frac{2}{2}$ as of August 6, 2021, 7 pm EST (by age - 10 year groups)



Date of illness onset

Figure 3. COVID-19 cases (n=1,434,642 1) in Canada by date of illness onset 2 as of August 6, 2021, 7 pm EST (by age - 20 year groups)




This figure may underestimate the total number of cases among returning travelers. Exposure history is not available for all cases and jurisdictions have not all consistently reported exposure history to PHAC throughout the pandemic.


Demographics

We have detailed case report data from 1,435,035 cases. We know the age of patients in 100.00% of cases, and both age and gender in 99.70% of cases.

Of the cases reported in Canada so far, 50.3% were female and 35.7% were between 20 and 39 years old (Figure 4).

Figure 4. Age $\stackrel{\checkmark}{}$ distribution of COVID-19 cases (n=1,435,035 $\stackrel{1}{}$) in Canada as of August 6, 2021, 7 pm EST $\stackrel{4}{}$





Age by gender $\frac{4}{}$ distribution of COVID-19 cases (n=1,435,035 $\frac{1}{}$) in Canada, August 6, 2021, 7 pm EST

| Age group (years) | Number of cases with case reports (percentage) | Number of male cases (percentage) | Number of female cases (percentage) | Number of other cases (percentage) |
|-------------------------|--|-----------------------------------|-------------------------------------|------------------------------------|
| ≤19 | 277,301 (19.3%) | 143,094 (20.1%) | 134,197 (18.7%) | 10 (17.9%) |
| 20-29 | 275,604 (19.2%) | 141,726 (19.9%) | 133,863 (18.6%) | 15 (26.8%) |
| 30-39 | 235,535 (16.4%) | 117,355 (16.5%) | 118,169 (16.4%) | 11 (19.6%) |
| 40-49 | 209,584 (14.6%) | 101,307 (14.2%) | 108,266 (15.1%) | 11 (19.6%) |
| 50-59 | 186,145 (13.0%) | 92,795 (13.0%) | 93,345 (13.0%) | 5 (8.9%) |
| 60-69 | 114,974 (8.0%) | 59,855 (8.4%) | 55,116 (7.7%) | 3 (5.4%) |
| 70-79 | 60,340 (4.2%) | 29,936 (4.2%) | 30,403 (4.2%) | 1 (1.8%) |
| 80+ | 71,210 (5.0%) | 25,248 (3.5%) | 45,962 (6.4%) | 0 (0.0%) |
| Total | 1,430,693 (100%) | 711,316 (100%) | 719,321 (100%) | 56 (100%) |

How people were exposed $\frac{3}{2}$

In Canada , detailed case report data were provided for 1,435,035 cases. We have exposure history for 1,070,583 (74.6%) cases. The probable exposure setting of these cases ¹/₂ are:

- any exposure that occurred in Canada: 982,297 (91.8%), including
 - o from contact with a known COVID case: 595,122 (55.6%)
 - from contact with a traveller: 8,629 (0.8%)
 - from an unknown source: 378,546 (35.4%)
- currently unknown (information pending): 78,154 (7.3%)
- travelled outside of Canada: 10,132 (0.9%)

Cases following vaccination

Data extracted on August 05, 2021 for cases from December 14, 2020 up until July 17, 2021.

While the COVID-19 vaccines are effective, there is still a small percentage of the population who are vaccinated that will still be infected with COVID-19 if they are exposed to the virus that causes it. This means that even with high vaccine effectiveness, a small percentage of the population who are vaccinated against COVID-19 will still get sick and some may be hospitalized or even die as a result of their illness. It is also possible that a person could be infected just before or just after vaccination and still get sick. It typically takes about 2 weeks for the body to build protection after vaccination, so a person could get sick if the vaccine has not had enough time to provide protection.


Cases reported since the start of the vaccination campaign, as of July 17, 2021

Since the start of the vaccination campaign on December 14, 2020, PHAC received case-level vaccine history data for 75.3% (n=617,042) of COVID-19 cases aged 12 years or older.

Of these cases:

- 552,688 (89.6%) were unvaccinated at the time of their episode date
- 32,751 (5.3%) were not yet protected by the vaccine, as their episode date occurred less than 14 days after their first dose
- 28,484 (4.6%) were only partially protected by the vaccine, as their episode date occurred either 14 days or more after their first dose or less than 14 days after their second dose
- 3,119 (0.5%) were fully protected by the vaccine, as their episode date occurred 14 days or more after their second dose

Figure 5. Distribution of confirmed COVID-19 cases reported to PHAC by vaccination status as of July 17, 2021



fully vaccinated confirmed cases reported to PHAC, as of July 17, 2021

| Status | Cases | Hospitalizations | Deaths |
|-------------------------|-------|------------------|--------|
| Unvaccinated | 89.6% | 84.8% | 82.1% |
| Cases not yet protected | 5.3% | 7.4% | 8.9% |
| Partially protected | 4.6% | 7.1% | 7.8% |
| Fully protected | 0.5% | 0.7% | 1.2% |

Among the eleven jurisdictions currently reporting case-level vaccine history data to PHAC, a total of 19.6 million people have received at least one dose of the COVID-19 vaccine as of the COVID-19 vaccine as of July 17, 2021.

Of these people:

- 19.1 million achieved partial vaccination status, of which 28,484 (0.15%) were diagnosed with COVID-19 while partially vaccinated
- 10.4 million achieved full vaccination status, of which 3,119 (0.03%) were diagnosed with COVID-19 while fully vaccinated

Based on detailed case information reported to PHAC from provinces and territories, cases following vaccination were reported more frequently among females and those aged 60 years and older (Table 2). This may be the result of higher vaccination coverage in Canada among females and those aged 60 years and older due to the prioritization of older age groups and healthcare workers as part of the vaccine rollout.

Table 2. Characteristics and severe outcomes associated unvaccinated, partially vaccinated and fully vaccinated confirmed cases reported to PHAC, as of July 17, 2021

| | | Unvaccinated (n=552,688) | Cases not yet protected (n=32,751) | Partially protected (n=28,484) | Fully protected (n=3,119) | Total cases (n=617,042) |
|-----------|---------|--------------------------|------------------------------------|--------------------------------|---------------------------|----------------------------|
| Gender* | Male | 281,999 (90.6%) | 15,599 (5.0%) | 12,653 (4.1%) | 887 (0.3%) | 311,138 (100%) |
| Gender | Female | 268,051 (88.5%) | 17,036 (5.6%) | 15,731 (5.2%) | 2,218 (0.7%) | 303,036 (100%) |
| Hospitali | zations | 27,941 (84.8%) | 2,446 (7.4%) | 2,327 (7.1%) | 241 (0.7%) | 32,955 (100%) |
| Deaths | | 5,896 (82.1%) | 636 (8.9%) | 561 (7.8%) | 89 (1.2%) | 7,182 (100%) |

Source: Detailed case information received by PHAC from provinces and territories, since December 14, 2020 **Note**:

- There are currently eleven jurisdictions reporting case-level vaccine history data to PHAC as part of the national COVID-19 dataset. A data cutoff of July 17, 2021 was used to account for any reporting delays associated with vaccine history information.
- *Cases with missing gender were excluded. Where available, gender data was used; when gender data was unavailable, sex data was used. Reliable data on gender diverse respondents are unavailable due to small counts.

Fully vaccinated individuals diagnosed with COVID-19 were significantly protected from severe outcomes. Compared to unvaccinated cases, fully vaccinated cases were 70% less likely to be hospitalized and 51% less likely to die as a result of their illness (Table 3).

Table 3. Risk of severe outcomes among fully vaccinated cases compared to unvaccinated cases as of July 17, 2021

| Severe Outcome | Adjusted* Odds Ratio (95% CI) | | |
|------------------|-------------------------------|--|--|
| Hospitalizations | 0.30 (0.26 - 0.35) | | |
| Deaths | 0.49 (0.38 - 0.62) | | |

^{*}Adjusted for 10-year age groups and month of episode date

Source: Detailed case information received by PHAC from provinces and territories

Note: Due to the nature of the dataset (i.e. confirmed cases of COVID-19 in Canada), the odds of severe outcomes among cases following vaccination only considers vaccinated individuals that contracted COVID-19. It does not reflect the protection conferred by the vaccines to prevent COVID-19 infection.

Data for this analysis is extracted from the COVID-19 national data set, which contains detailed case-level information received by PHAC from all provinces and territories. Note that a data cut-off of July 17, 2021 was used to account for any reporting delays associated with vaccine history information. There are currently eleven jurisdictions reporting case-level vaccine history data to PHAC as part of the national COVID-19 dataset.

PHAC monitors cases following vaccination using the following categories:

Unvaccinated cases: include those who were unvaccinated at the time of their episode date.

Cases not yet protected from vaccination include those whose episode date occurred less than 14 days after their first dose of the vaccine.

Partially vaccinated cases include those whose episode date occurred 14 days or more after their first vaccine dose or less than 14 days after their second dose of the vaccine.

Fully vaccinated cases include those whose episode date occurred 14 days or more after their second dose of a two dose vaccine series or those whose episode date occurred 14 days or more after one dose of a one-dose


vaccine.

Note: When symptom onset date is unavailable or the case is asymptomatic, episode date uses the following dates as a proxy for classification: laboratory specimen collection date, or laboratory testing date.

Severe illness and outcomes

Hospital use

Figure 6. Daily number of hospital beds and ICU beds occupied by COVID-19 patients as of August 2, 2021



Between July 26, 2021 and August 2, 2021:

- the number of **hospital beds** occupied by COVID-19 patients **decreased** from **322** to **315** beds.
- the number of ICU beds occupied by COVID-19 patients decreased from 188 to 166 beds.
- the number of COVID-19 patients who were mechanically vented decreased from 118 to 109.

Hospitalizations and deaths to date

We have detailed case report data on 1,435,035 cases, and hospitalization status for 1,006,988 (70.2%) of them:

- 75,642 cases (7.5%) were hospitalized, of whom:
 - 14,355 (19.0%) were admitted to the ICU
 - 1,924 (2.5%) needed mechanical ventilation

The provinces and territories provided detailed case report forms for **26,646** deaths related to COVID-19.

Figure 7a. Age and gender $\frac{4}{}$ distribution of COVID-19 cases hospitalized in Canada as of August 6, 2021, 7 pm EST (n=75,500 $\frac{1}{}$)




Figure 7b. Age and gender 4 distribution of COVID-19 cases admitted to ICU in Canada as of August 6, 2021, 7 pm EST (n=14,197 1)





Figure 7c. Age and gender $\frac{4}{}$ distribution of COVID-19 cases deceased in Canada as of August 6, 2021, 7 pm EST (n=26,591 $\frac{1}{}$)



Data note: Figure 7 includes COVID-19 cases hospitalized, admitted to ICU, and deceased for which age and gender information were available. Therefore, some COVID-19 hospitalizations, ICU admissions, and deaths may not be included in Figure 7.

Age and gender $\frac{4}{}$ distribution of COVID-19 cases hospitalized in Canada as of August 6, 2021, 7 pm EST (n=75,500 $\frac{1}{}$)

| Age group (years) | Number of cases with case reports (percentage) | Number of male cases (percentage) | Number of female cases (percentage) | Number of other cases (percentage) |
|-------------------------|--|-----------------------------------|-------------------------------------|------------------------------------|
| 0-19 | 1,418 (1.9%) | 758 (1.0%) | 659 (0.9%) | 1 (0.0%) |
| 20-29 | 3,069 (4.1%) | 1,286 (1.7%) | 1,783 (2.4%) | 0 (0.0%) |
| 30-39 | 5,212 (6.9%) | 2,501 (3.3%) | 2,711 (3.6%) | 0 (0.0%) |
| 40-49 | 6,822 (9.0%) | 3,953 (5.2%) | 2,869 (3.8%) | 0 (0.0%) |
| 50-59 | 11,027 (14.6%) | 6,629 (8.8%) | 4,398 (5.8%) | 0 (0.0%) |
| 60-69 | 13,351 (17.7%) | 7,964 (10.5%) | 5,387 (7.1%) | 0 (0.0%) |
| 70-79 | 14,630 (19.4%) | 8,138 (10.8%) | 6,492 (8.6%) | 0 (0.0%) |
| 80+ | 19,971 (26.5%) | 9,149 (12.1%) | 10,822 (14.3%) | 0 (0.0%) |

Age and gender $\frac{4}{}$ distribution of COVID-19 cases admitted to ICU in Canada as of August 6, 2021, 7 pm EST (n=14,197 $\frac{1}{}$)

| Age group (years) | Number of cases with case reports (percentage) | Number of male cases (percentage) | Number of female cases (percentage) | Number of other cases (percentage) |
|-------------------------|--|-----------------------------------|-------------------------------------|------------------------------------|
| 0-19 | 165 (1.2%) | 89 (0.6%) | 76 (0.5%) | 0 (0.0%) |
| 20-29 | 384 (2.7%) | 199 (1.4%) | 185 (1.3%) | 0 (0.0%) |
| 30-39 | 854 (6.0%) | 495 (3.5%) | 359 (2.5%) | 0 (0.0%) |
| 40-49 | 1,448 (10.2%) | 907 (6.4%) | 541 (3.8%) | 0 (0.0%) |
| 50-59 | 2,827 (19.9%) | 1,860 (13.1%) | 967 (6.8%) | 0 (0.0%) |
| 60-69 | 3,777 (26.6%) | 2,438 (17.2%) | 1,339 (9.4%) | 0 (0.0%) |
| 70-79 | 3,270 (23.0%) | 2,080 (14.7%) | 1,190 (8.4%) | 0 (0.0%) |
| 80+ | 1,472 (10.4%) | 835 (5.9%) | 637 (4.5%) | 0 (0.0%) |

Age and gender 4 distribution of COVID-19 cases deceased in Canada as of August 6, 2021, 7 pm EST (n=26,591 1)

| Age group (years) | Number of cases with case reports (percentage) | Number of male cases (percentage) | Number of female cases (percentage) | Number of other cases (percentage) |
|-------------------------|--|-----------------------------------|-------------------------------------|------------------------------------|
| 0-19 | 15 (0.1%) | 6 (0.0%) | 9 (0.0%) | 0 (0.0%) |
| 20-29 | 66 (0.2%) | 42 (0.2%) | 24 (0.1%) | 0 (0.0%) |
| 30-39 | 148 (0.6%) | 96 (0.4%) | 52 (0.2%) | 0 (0.0%) |
| 40-49 | 344 (1.3%) | 225 (0.8%) | 119 (0.4%) | 0 (0.0%) |
| 50-59 | 1,001 (3.8%) | 621 (2.3%) | 380 (1.4%) | 0 (0.0%) |
| 60-69 | 2,573 (9.7%) | 1,646 (6.2%) | 927 (3.5%) | 0 (0.0%) |
| 70-79 | 5,397 (20.3%) | 3,229 (12.1%) | 2,168 (8.2%) | 0 (0.0%) |
| 80+ | 17,047 (64.1%) | 7,547 (28.4%) | 9,500 (35.7%) | 0 (0) |

Provincial, territorial and international reporting

For more information, please refer to provincial or territorial COVID-19 webpages:

- British Columbia
- Alberta
- Saskatchewan
- Manitoba
- Ontario
- Quebec
- · Newfoundland and Labrador
- New Brunswick
- Nova Scotia
- Prince Edward Island
- Yukon
- Northwest Territories
- Nunavut
- World Health Organization
- Centers for Disease Control and Prevention
- European Centre for Disease Control and Prevention
- This figure is based on cases for which a case report form was received by the Public Health Agency of Canada from provincial or territorial partners.
- The shaded area represents a period of accumulating data where it is expected that cases have occurred but have not yet been reported nationally. The earliest of the following dates were used as an estimate: Onset date, Specimen Collection Date, Laboratory Testing Date, Date Reported to Province or Territory, or Date Reported to PHAC.
- Exposure information may not be available for all cases. Some jurisdictions haven't consistently reported to PHAC how people were exposed throughout the pandemic. As a result, this may underestimate the total number of cases by different exposures, especially among returning travelers.
- Where available, gender data was used; when gender data was unavailable, sex data was used. Reliable data on gender diverse respondents are unavailable due to small counts.